

The Expression of Genes Encoding Secreted Proteins in *Medicago truncatula* A17 Inoculated Roots

LUCIA KUSUMAWATI^{1,2*}, KATHRYN KURAN², NIJAT IMIN²,
ULRIKE MATHESIUS², MICHAEL DJORDJEVIC²

¹Department of Biotechnology, Faculty of Life Sciences, Surya University,
Jalan Scientia Boulevard blok U/7, Gading Serpong, Tangerang 15810, Indonesia

²Research School of Biology, The Australian National University, Acton ACT 0200, Australia

Received February 12, 2013/Accepted August 2, 2013

Subtilisin-like serine protease (MtSBT), serine carboxypeptidase (MtSCP), MtN5, non-specific lipid transfer protein (MtnsLTP), early nodulin2-like protein (MtENOD2-like), FAD-binding domain containing protein (MtFAD-BP1), and rhicadhesin receptor protein (MtRHRE1) were among 34 proteins found in the supernatant of *M. truncatula* 2HA and *sickle* cell suspension cultures. This study investigated the expression of genes encoding those proteins in roots and developing nodules. Two methods were used: quantitative real time RT-PCR and gene expression analysis (with promoter:*GUS* fusion) in roots. Those proteins are predicted as secreted proteins which is indirectly supported by the findings that promoter:*GUS* fusions of six of the seven genes encoding secreted proteins were strongly expressed in the vascular bundle of transgenic hairy roots. All six genes have expressed in 14-day old nodule. The expression levels of the selected seven genes were quantified in *Sinorhizobium*-inoculated and control plants using quantitative real time RT-PCR. In conclusion, among seven genes encoding secreted proteins analyzed, the expression level of only one gene, *MtN5*, was up-regulated significantly in inoculated root segments compared to controls. The expression of *MtSBT1*, *MtSCP1*, *MtnsLTP*, *MtFAD-BP1*, *MtRHRE1* and *MtN5* were higher in root tip than in other tissues examined.

Keywords: secreted proteins, *Medicago truncatula*, root, nodules, RT-PCR, gene expression

INTRODUCTION

To adapt to low nitrogen availability in soil, some plants form a symbiosis with nitrogen fixing bacteria to provide a nitrogen source for the plants. This symbiosis between legumes and rhizobia give rise to a completely new organ called nodule. Nodules (and lateral roots) are initiated from differentiated root cells that become meristematic. In nodules, those meristematic cells will differentiate into different nodule tissues. Nodule formation involves signal exchange and recognition, cell differentiation and cell division that eventually lead to the formation of a completely new organ.

The symbiosis between legume plants and rhizobia involves various physiological processes performed by plants to adapt to their changing environment. This includes signal exchange and recognition, regulation of defence responses and re-initiation of cell divisions that eventually lead to the formation of a completely new organ, a root nodule (Oldroyd & Downie 2008). The symbiotic interaction shows a high degree of host-rhizobia specificity. It is started when roots of

host plants secrete flavonoids or isoflavonoids that attract rhizobia to colonize the rhizosphere and induce the rhizobia nodulation (nod) genes (Gulash *et al.* 1984; Peters *et al.* 1986; Redmond *et al.* 1986; Kape *et al.* 1991). Each rhizobial species is activated by a particular set of flavonoids. The induction of nod genes results in the secretion of the nodulation factors (Nod factors), lipo-chitin oligosaccharide molecules with variable structures. Different rhizobia produce different Nod factors that determine the host range of the bacteria (Roche *et al.* 1991; Ehrhardt *et al.* 1995; Lorquin *et al.* 1997). Once the rhizobia attach themselves to the root hairs, the Nod factors induce root hair curling and further distortion into a 'shepherd's crook'. Within the pocket which has been formed by the curled hair, bacterial intrusion takes place and is facilitated by the development of an infection thread (Gage & Margolin 2000). Nod factors also induce cortical cell division of host plant, coinciding with the induction of several specific genes in early nodulation (early nodulin genes) (Bisseling *et al.* 1990).

Nodule development is affected by phytohormones (Scheres *et al.* 1992; Nukui *et al.* 2000; Wasson *et al.* 2006), defence response of the host plant (Niehaus

*Corresponding author. Phone: +62-21-70207727,
Fax: +62-21-70200722, E-mail: lucia.kusumawati@surya.ac.id

et al. 1993; Perotto *et al.* 1994; Martínez-Abarca *et al.* 1998) and the physiological condition of the host plant (Streeter & Wong 1988; Caetano-Anolles & Gresshoff 1991).

Generally, secreted proteins are defined as proteins that are transported intracellularly through the endomembrane system and then delivered to the extracellular environment by means of secretory vesicles that fuse with the cell membrane (Jones & Robinson 1989). In the extracellular environment, secreted proteins can diffuse away and some are freely soluble in the apoplastic space (apoplastic proteins). They also can become attached to the cell wall or become incorporated into the cell membrane where they act as membrane transporters or ion channels, ligand receptors and signalling complexes, or contact points for intracellular cytoskeletal or for the extracellular matrix (Greenbaum *et al.* 2001; Sanderfoot & Raikhel 2003; Lee *et al.* 2004).

Previous studies have revealed that secreted proteins are involved in pathogenic stresses (Rep *et al.* 2002; Grunwald *et al.* 2003; Ndimba *et al.* 2003; Hugot *et al.* 2004), environmental stresses (Fecht-Christoffers *et al.* 2003), cell-cell recognition (Franklin-Tong & Franklin 2003) and development (Pearce *et al.* 2001; von Groll *et al.* 2002; Casamitjana-Martinez *et al.* 2003). In nodule development, physiological processes such as defense response, meristematic cell division and differentiation are taken place.

Examples of proteins involved in the maintenance of shoot apical meristem (SAM) are CLV1 (a receptor like kinase with a leucine rich repeat extracellular domain), CLV2 (a receptor like protein with leucine rich repeat extracellular domain but without the kinase cytoplasmic domain), CLV3 (a secreted protein that acts as a ligand for CLV1) (Clark *et al.* 1997; Jeong *et al.* 1999; Fletcher *et al.* 1999). Secreted protein phyto-sulfokine- α from *Asparagus officinalis* (Matsubayashi & Sakagami 1996) and *Oryza sativa* (Matsubayashi *et al.* 1997) is involved in promoting cell proliferation and inducing cell differentiation in *Zinnia elegans* (Matsubayashi *et al.* 1999). It is been known that genes encoding receptor-like kinase (RLK) protein, when mutated, cause excessive number of nodules. They are *har1* (from *Lotus japonicus*), *nts1* (from *Glycine max*), *sym29* (from *Pisum sativum*), and *sun* (from *Medicago truncatula*) (Wopereis *et al.* 2000; Krusell *et al.* 2002; Nishimura *et al.* 2002; Searle *et al.* 2003; Schnabel *et al.* 2005). However, little is known about the potential roles and identities of secreted proteins during the *Rhizobium*-legume symbiosis. Proteome analysis of soybean xylem sap

identified a xyloglucan transendoglycosyl transferase was upregulated 8 h after inoculation with *Bradyrhizobium japonicum*. However, silencing of the gene encoding this secreted protein did not result in changes in nodulation (Subramanian *et al.* 2009). Overall, it is likely that secreted proteins play a role in the regulation of meristem initiation and control of defense responses during nodulation.

Here, we used quantitative real time RT-PCR to study the expression level of genes encoding several secreted proteins: subtilisin-like serine protease (*MtSBT*), serine carboxypeptidase (*MtSCP*), *MtN5*, non-specific lipid transfer protein (*MtnsLTP*), early nodulin2-like protein (*MtENOD2-like*), FAD-binding domain containing protein (*MtFAD-BP1*), and rhicadhesin receptor protein (*MtRHRE1*), in *M. truncatula* roots and nodules. Their expression in roots and nodules were also compared to other organs (leaves and cotyledons). Previously, quantitative real time RT-PCR has been used to study the expression level of genes (Prayitno *et al.* 2006a; Holmes *et al.* 2008). Further, this study also studied the expression of those genes in root system and nodules using *GUS* reporter gene assay.

MATERIALS AND METHODS

All chemicals were from Sigma-Aldrich (St. Louis, MO) unless stated otherwise. Deionized water 18.2 M Ω /cm ("MQ water") was used throughout the study.

***Medicago truncatula* A17 growth.** *Medicago truncatula* A17 seeds were grown as described in Prayitno *et al.* (2006b). *M. truncatula* seeds were scarified and then surface sterilized with 6.25% (v/v) sodium hypochlorite for 15 minutes. After seven washes with sterile water, seeds were spread on nitrogen-free Fåhræus agar media (Fåhræus 1957) containing 0.8% agar (Agar Grade J3, Gelita Pty Ltd., Beaudesert, Queensland, Australia). Seeds were incubated in the dark at 4 °C for one day to break their dormancy and were then germinated by incubating the plate upside down in the dark at 28 °C overnight, so that emerging roots grew straight into the air. Seedlings with similar root length were selected and transferred to 15 cm diameter Petri dishes containing nitrogen-free Fåhræus agar media and sealed $\frac{3}{4}$ around with parafilm. The seedlings were incubated vertically in the growth chamber with a photon flux density of 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for 16 h at 20 °C for four days. To reduce light intensity around the roots, pieces of thick-dark paper were placed in the lower half of the plate between the plates. Under these conditions, seedlings grew on the media

surface, and were easy to inoculate and to observe under sterile conditions. After four days incubation, the seedlings were flood-inoculated at 1 cm above the root tip with 5 μ L of diluted *Sinorhizobium meliloti* strain 1021 culture (at OD₆₀₀ of 0.1). Two weeks after inoculation, tissue samples were taken from leaves, cotyledon, root segments, and root tips.

Agrobacterium rhizogenes–Mediated Hairy Root Transformation. *Medicago truncatula* A17 seeds were scarified and surface-sterilised with 6.25% (v/v) sodium hypochlorite for 15 minutes. After seven washes with sterile water, seeds were grown as described in Prayitno *et al.* (2006b). The transformation of *M. truncatula* with *A. rhizogenes* was done as described in Boisson-Dernier *et al.* (2001).

Bacterial Growth. *Sinorhizobium meliloti* strain 1021 was grown in Bergensen's Modified Medium (BMM) agar media (Rolfe *et al.* 1980) for 2 days. A colony of bacteria was then transferred to BMM liquid medium and grown at 28 °C overnight. The culture was then diluted with sterile water to an OD₆₀₀ of 0.1 for inoculation of plants. *Agrobacterium rhizogenes* Arqual strain (Boisson-Dernier *et al.* 2001) was maintained on Luria-Bertani (LB) agar plates containing 100 μ g mL⁻¹ streptomycin.

DNA Isolation. Genomic DNA was isolated using a PUREGENE™ DNA purification kit according to the manufacturer's instructions (Gentra System, Minneapolis, Minn, USA).

Primer Design Promoter Primers. The open reading frame from Softberry was blasted to the University of Oklahoma *Medicago truncatula* genome blast server (http://www.genome.ou.edu/medicago_blast.html) using blastn to get the contig number (for example mtgsp_008f03.Contig1 for FAD). BAC sequences were downloaded from the University of Oklahoma (<ftp://ftp.genome.ou.edu/pub/medicago>) and the previously obtained contig number was used to get the complete sequence of the relevant gene. At least two Kb of upstream sequence from the start codon was selected and primers were designed using the online application from Integrated DNA Technologies (<http://www.idtdna.com/analyzer/Applications/OligoAnalyzer/Default.aspx>). Primers were chosen to have a GC content between 40% to 60%, melting temperature above 55 °C and a minimum chance to allow a self dimer of < 8 to form. The oligos were also checked to make sure that they do not have a strong hairpin structure. Primers used for promoter cloning were: SUBT_Promf 5'-CACCCGCTGAGTTAAGACAAGGTTTGTG-3', SUBT_Promr 5'-TCTCACCTCTTGATTTTGG AATGAG-3', FAD_Promf 5'-CACCTCTATAA

GTTGGGCGAGGGATA-3', FAD_Promr 5'-GA TTTTAGTTTGTGTTGTGTTTGTGTTTGT-3', Serine_Promf 5'-CACCAGTTGAAGGAATGG CCGG-3', Serine_Promr 5'-TGCTGAAGAGGAA GCAGAAAC-3', nsLTP_Promf 5'-CACCTGCGTA TGCGGGAGCTT-3', nsLTP_Promr 5'-TCACCAA TAATTTTGTGTTGTA CTTC AATAAT-3', MtN5_Promf 5'-CACCGCGAGCCAACATCAA CTAATTTTTC-3', MtN5_Promr 5'-ACTCCTAA TTAATTAACCA AATGCTTAGCT-3', Enod2_Promf 5'-CACCAAGTTCATCAAAAGC ATGTCATCAGA-3', Enod2_Promr 5'-TGTTAATT GCTAAATATTTAGAGAAACA-3', Rhic_Promf 5'-CACCGCCAAAGACGCCATAA TGATC-3', Rhic_Promr 5'-GTTAAGATCGGTGAC AAACAAATTAACC-3'

Quantitative Real Time RT-PCR Primers.

Gene-specific primers were designed using Primer Express software (Applied Biosystems, Foster City, CA, USA) and synthesized by GeneWorks (Adelaide, Australia). Those primer were MtSBT1_F 5'-GC TTCAACGATGCTGGGTTT-3', MtSBT1_R 5'-CTGATTCACACGCGCCTTT-3', MtSCP1_F 5'-CCA ACTTGCA CA ACTCATTGTTTC-3', MtSCP1_R 5'-TCCCTATTGCAATTCCTTGA-3', MtN5_F 5'-AGGGCAGATGAGGCTTGCT-3', MtN5_R 5'-CGCAGAGGCAAGGAAGATTG-3', MtnsLTP_F 5'-TTTGCTGCTCGGCTATTGC-3', MtnsLTP_R 5'-GCAATCCTGAATCCTTATAA CGACAT-3', MtENOD2-like_F 5'-CAAACCC AATCAAGAAGTTCGAA-3', MtENOD2-like_R 5'-GGCCCTGCTCTGTCCAAAG-3', MtFAD-BP1_F 5'-GGCATCGGTGGCCTCAT-3', MtFAD-BP1_R 5'-CCAAGGCCATACTTTCTCATCAT-3', MtRHRE1_F 5'-TCATTGGCACGTATTGACTA TGC-3', MtRHRE1_R 5'-TGCACGTGGGTGAG TGTGA-3'

Polymerase Chain Reactions. Polymerase Chain Reactions (PCR) were performed in Thermo Hyaid PCR Express Thermal Cycler (Fisher Scientific, Pittsburgh, PA, USA) using Phusion Hot Start High-Fidelity DNA Polymerase (Finnzymes, Finland). The initial denaturations were carried out at 98 °C for 30 seconds followed by a total of 35 cycles as follows: denaturation at 98 °C for 10 seconds, annealing between 55 °C and 65 °C for 30 seconds, and extension at 72 °C for 1 min/kb. The final extension was carried out at 72 °C for 5 minutes.

Gateway Cloning. The GATEWAY™ Cloning Technology was used to generate expression vectors incorporating entry vectors. PCR products were cloned into an entry vectors using pENTR™ Directional TOPO Cloning Kit (Invitrogen, Carlsbad, CA, USA) to create a Gateway® entry clones. TOPO

Cloning reactions were transformed into Premade Z-Competent™ *E. coli* cells (Zymo Research, CA, USA) following the manufacturer's instructions. Plasmid DNA (entry clones) were isolated using QIAprep Spin Miniprep kit (QIAGEN, Melbourne, Australia) and entry clones were confirmed by PCR and gel electrophoresis.

The sequences of the entry constructs were confirmed by DNA sequencing at the Biomolecular Resource Facility, The John Curtin School of Medical Research, the Australian National University. Subsequently, correct sequences of interest were transferred from the entry clones into Gateway® destination vectors to generate expression clones by performing the LR recombination reaction using Gateway® LR Clonase™ II enzyme mix (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions.

LR reaction mixtures were transformed into Premade Z-Competent™ *E. coli* cells (Zymo Research, CA, USA) as described in the manufacturer's protocol and the cells were grown overnight on LB plates containing 50 µg mL⁻¹ spectinomycin and 100 µg mL⁻¹ streptomycin. Putative positive clones (transformed *E. coli* containing expression clones) were cultured overnight in liquid LB containing the same antibiotics. Plasmid DNA (expression clones) were isolated using QIAprep Spin Miniprep kit (QIAGEN, Melbourne, Australia) and expression clones were confirmed by PCR and gel electrophoresis. One µg of expression clones DNA was transformed into 100 µL *Agrobacterium rhizogenes* ARqual strain (Boisson-Dernier *et al.* 2001) using the freeze-thaw transformation method (Höfgen & Willmitzer 1988). Transformants were selected by resistance to spectinomycin and streptomycin.

β-Glucuronidase (GUS) Staining. GUS activity was localized in whole transgenic root of *M. truncatula* following the protocol described previously (Vitha *et al.* 1995).

Statistical Analysis. Student's *t* test was performed using Microsoft Excel 2003.

Gene Expression Analysis. The expression levels of the seven selected genes were examined by using relative transcript expression which was obtained from quantitative real-time RT-PCR. The examined tissues were root segments (approximately 2 cm segments where nodules usually formed), root tips (0.5 cm from the tip), leaves and cotyledons. Each tissue was divided into two groups, i.e. non-inoculated (control) and inoculated with *Sinorhizobium liloti*. Samples were harvested 14-day post inoculation with consideration that it is quite difficult

to differentiate nodules and lateral root primordia before 14 day. Quantitative real-time RT-PCR data was analyzed using Student's *t* test.

RESULTS

Gene Expression Analysis. As shown in Figure 1, the expression level of subtilisin-like serine protease (*MtSBT1*), serine carboxypeptidase (*MtSCPI*), *MtN5*, non-specific lipid transfer protein (*MtnsLTP*), FAD-binding domain containing protein (*MtFAD-BP1*) and rhicadhesin receptor (*MtRHRE1*) were higher in root tips compared to its expression in other tissues (leaves, cotyledons and root segments). Only early nodulin 2-like (*MtENOD2-like*) was expressed higher in leaves than in roots. The high expression level of *MtN5* in root tips (Figure 1C) was also reported previously by Holmes *et al.* (2008). Interestingly, as shown by Figure 1C, *MtN5* expression was up-regulated 15-fold ($P < 0.05$) in inoculated root segments compared to control root segments (non-inoculated). There was no significant difference in *MtN5* expression between inoculated leaves, cotyledons, root tips and their control (non-inoculated) counterparts (Figure 1C). There were also no significant difference in the expression level of *MtSBT1*, *MtSCPI*, *MtnsLTP*, *MtFAD-BP1*, and *MtRHRE1* between control (non-inoculated) and inoculated group for any tissue examined.

The location of the expression of genes encoding six secreted proteins in roots was also examined using *A. rhizogenes*-mediated hairy root transformation. Six promoters were successfully isolated and cloned while the promoter of *MtN5* could not be isolated successfully after several attempts. Therefore, the location of *MtN5* expression in roots and nodules could not be reported here.

In transformed roots, *MtSBT1:GUS* transgenic roots showed strong expression in the vascular bundle and root apex (Figure 2A). At different stages of lateral root development, *MtSBT1:GUS* was expressed stronger in the zone where lateral roots are initiated (Figure 2A). In nodules, it was expressed in all areas of the nodule but stronger in the vascular bundle, apex and basal nodule region (Figure 3A). Figure 2B demonstrates the expression of *MtSCPI:GUS* in transgenic roots, which showed expression in the vascular bundle, in both roots and nodules. It was also expressed strongly in the apex zone of root tips and where lateral roots were initiated, confirming the RT-PCR data that *MtSCPI* was expressed higher in root tips than other tissues.

In transformed roots, *MtnsLTP:GUS* did not seem to be expressed in the main root, except in the pex




Figure 1. The expression level of genes encoding seven predicted secreted proteins. Gene expression level of (A) *MtSBT1*, (B) *MtSCP1*, (C) *MtN5*, (D) *MtnsLTP*, (E) *MtENOD2-like*, (F) *MtFAD-BP1*, (G) *MtRHRE1*. Relative transcript expression is obtained by mean of real-time RT-PCR. Sample used was *Medicago truncatula* A17 which was divided into two sample groups, i.e. control – non-inoculated plants (Ctrl) and 14 day post inoculated plants (Inoc). Each sample group presents leaf, cotyledon (Cotyl), root segment and root tip. Error bars show standard deviations.




Figure 2. *GUS* reporter gene assay for transgenic root of *M. truncatula* A17. The expression of *GUS* reporter gene in transgenic root driven by promoter of (A) *MtSBT1*, (B) *MtSCP1*, (C) *MtnsLTP*, (D) *MtENOD2-like*, (E) *MtFAD-BP1*, (F) *MtRHRE1*. *GUS* expression was observed in different stages of lateral root development (L), root tip (T), and vascular bundle (V). White bars represent 0.5 mm.

zone and the first centimeter of root tip (Figure 2C). Nonetheless, its expression in some of the lateral roots was detected, not limited to the vascular bundle. During different developmental stages of the lateral root, *MtnsLTP:GUS* expression was localized in the apex of root tip and where lateral roots were initiated (Figure 2C). In inoculated roots, *MtnsLTP:GUS* was expressed in the vascular bundle and in the entire

nodule (Figure 3C). Its expression was stronger in apex. In general, there was a parallel between RT-PCR data and *MtnsLTP:GUS* expression localization. RT-PCR data showed that *MtnsLTP* expressed the highest in root tips (Figure 1D) and *MtnsLTP:GUS* showed a high expression in root apex and the first centimetre from the tip (Figure 2C).




Figure 3. GUS staining of nodules formed in transgenic roots expressing promoter:*GUS* construct. The expression of *GUS* reporter gene in transgenic root driven by promoter of (A) *MtSBT1*, (B) *MtSCP1*, (C) *MtnsLTP*, (D) *MtENOD2-like*, (E) *MtFAD-BP1*, (F) *MtRHRE1*. White bars represent 0.5 mm.

In transformed roots, *MtENOD2-like:GUS* expression was mainly in the vascular tissue (Figure 2D). It was also expressed in the apex, either in the root tip or in the location where lateral root primordial is initiated. In 14-day old nodules, it was expressed in apex and vascular tissue (Figure 3D). RT-PCR data showed that *MtENOD2-like* was expressed higher in aerial organs than in root segments and inoculated root segments (Figure 1E). *MtENOD2-like:GUS* expression in transformed roots also showed a relatively low expression in 14-day old nodules (Figure 3D).

As can be seen in Figure 2E, *GUS* reporter gene (under *MtFAD-BP1* promoter) was expressed strongly in vascular tissues and apex. During lateral root development, it was expressed in dividing cells where the growth was initiated and in vascular tissue. In nodules, it was expressed in vascular tissue, in the apex and base of the nodule (Figure 3E).

In Figure 2F, *GUS* reporter gene (under *MtRHRE1* promoter) was expressed in vascular tissue and the root apex, either in root tips or in lateral root primordia. In 14-day old nodules, it was expressed in the entire nodule but stronger in nodule apex (Figure 3F).

DISCUSSION

The expression of selected genes encoding secreted proteins was studied using two different dataset, i.e. *GUS* expression in transformant roots and quantitative real time RT-PCR data.

GUS Expression in Transformant Roots. In all transgenic roots, *GUS* expression was detected in nodules (Figure 3). It was also detected in the dividing and enlarging cells where the lateral roots were initiated in root apex and in vascular bundle (Figure 2). The expression of *GUS* in actively proliferating cells suggests that these secreted proteins have role in cell division and growth. Further, the use of promoter:*GUS* constructs showed that all six promoters examined were strongly expressed in vascular bundles (Figure 2 & 3). They were *MtSBT1*, *MtSCP1*, *MtnsLTP*, *MtFAD-BP1*, *MtRHRE1*, and *MtENOD-like2*. Their strong expression in the vascular bundle suggests that they might be involved in vascular bundle development or function, and that the gene products would be released into apoplast on a continuous basis along the entire length of the vasculature.

It is interesting to note that transgenic roots carrying *MtSBT1:GUS*, *MtnsLTP:GUS*, and *MtRHRE1:GUS* were expressed strongly the whole nodule. Therefore, it is possible that those secreted proteins have a role in nodule development.

Expression Level of *MtSBT1*, *MtSCP1*, *MtN5*, *MtnsLTP*, *MtFAD-BP1* and *MtRHRE1*. Using quantitative real time RT-PCR, six genes encoding secreted proteins examined showed higher expression in root tips than in leaves, cotyledons or root segments. These six genes were *MtSBT1*, *MtSCP1*, *MtN5*, *MtnsLTP*, *MtFAD-BP1* and *MtRHRE1*. Only *MtENOD2-like* showed lower expression in roots (either inoculated or non-inoculated) than in leaves and cotyledons (Figure 1).

Subtilisin-Like Serine Protease (*MtSBT1*) and Serine Carboxypeptidase (*MtSCP1*). In *Alnus glutinosa*, a subtilase, *Ag12*, was detected in nodules but not in roots, cotyledons, flowers and developing fruits (Ribeiro *et al.* 1995). In *Casuarina glauca*, *Cg12* (a *Ag12* homolog) was also expressed in nodules but not in roots, stems and leaves (Laplaze *et al.* 2000). Both were expressed stronger in the young infected cells of the infection zone, weaker in the fixation zone and no expression of *Ag12* and *Cg12* was detected in senescent infected cells (Ribeiro *et al.* 1995; Laplaze *et al.* 2000). Since young infected cells are active in synthesizing proteins, therefore, a stronger expression of *Cg12* and *Ag12* in the young infected cells suggests that *Cg12* and *Ag12* are involved in protein processing rather than proteolysis (Ribeiro *et al.* 1995; Laplaze *et al.* 2000). In this study, *MtSBT1* was expressed strongly in nodule apex, root apex and dividing cells. Therefore, it is possible that *MtSBT1* may have a role in cell division and growth in *M. truncatula*.

In *A. thaliana*, brassinosteroid insensitive 1 (*BRI1*) encodes a leucine-rich repeat receptor serine/threonine kinase (Li & Chory 1997; Friedrichsen *et al.* 2000) cell surface receptor for brassinosteroids, a plant growth and development regulator (He *et al.* 2000). A suppressor of weak *bri1* allele, *BRS1* (brassinosteroid insensitive 1 suppressor) was found by a gain-of-function genetic screen (Li *et al.* 2001). Using *35S:BRS1-GFP* transgenic plants, it was shown that *BRS1* (brassinosteroid insensitive 1) encodes a secreted serine carboxypeptidase (Zhou & Li 2005). Overexpression of *BRS1* in wild-type plants showed no obvious phenotype alteration while overexpression of *BRS1* in *bri1-5* and *bri1-9* mutants resulted in the partially restored phenotype caused by the point mutation in extracellular domain of *BRI1* by 52 and 75%, respectively (Noguchi *et al.* 1999; Friedrichsen *et al.* 2000; Li *et al.* 2001). In addition, overexpressing *BRS1* in a cytoplasmic kinase domain mutant, *bri1-1*, failed to restore the phenotype (Li *et al.* 2001). These observations suggest that *BRS1*, a secreted serine carboxypeptidase, involves in early events of *BRI1* signaling in the apoplast.

In this study, transgenic roots expressing *GUS* under *MtSCP1* promoter showed that, in mature nodules, *GUS* was expressed only in vascular bundle (Figure 3B), suggesting that *MtSCP1* involved in vascular bundle development or function.

Early Nodulin 2-Like Protein (*MtENOD2-like*), FAD-Binding Domain Containing Protein (*MtFAD-BP1*) and Rhicadhesin Receptor (*MtRHRE1*). Previously, *ENOD2* is known as a nodulin gene which is expressed specifically in

nodules (Gloude-mans & Bisseling 1989), and is used as a molecular marker for the early stage of nodule formation. Previous studies on nodulins compared their expression in nodules to their expression in roots but not to their expression in other organs (Fuller *et al.* 1983; Govers *et al.* 1985; Lang-Unnasch & Ausubel 1985), or compared nodulin encoded proteins to *Rhizobium* encoded proteins (Fuller *et al.* 1983; Lang-Unnasch & Ausubel 1985). However, the expression of *ENOD2* has been detected in uninfected roots of *Lotus japonicus* (Szczyglowski *et al.* 1997), *Glomus intraradices* colonised *M. sativa* roots (van Rhijn *et al.* 1997) and in cytokinin-treated *M. sativa* roots (van Rhijn *et al.* 1997). Further, *ENOD2* expression was also detected in the parenchyma cells of alfalfa bacteria free-nodules (van de Wiel *et al.* 1990). Additionally, *ENOD2-like* expression was detected in the stems, flowers and roots of non-nodulating tree legumes, *Styphnolobium japonicum* and *Cladrastis kentukea* (Foster *et al.* 2000). Due to the growing evidence that what was first thought to be nodule specific genes are also expressed in other parts of different plants, it has been proposed that genes involved in nodulation, such as *ENOD2*, are evolved from genes involved in general physiological processes which are common to most plants (Gualtieri & Bisseling 2000).

In this study, it was shown by quantitative real time RT-PCR that *MtENOD2-like* was expressed higher in leaves and cotyledons than in roots. Therefore, it is possible that *MtENOD2-like* may not have a specific role in root growth.

In transgenic roots that express reporter gene under *MtENOD2-like* promoter, *GUS* was expressed in the vascular tissue, in the apex of root tips and in the place where lateral roots are initiated. In 14-day old nodule, *GUS* expression was detected in vascular bundle at the base of the nodule and a very weak expression was detected at the apex of the nodule. A similar pattern was shown by transgenic roots that express *GUS* reporter gene under *MtFAD-BP1* promoter. In those roots, *GUS* was expressed very strongly in vascular bundle, in the apex of root tips and in the lateral root primordium. In 14-day old nodule, *GUS* was expressed in the vascular bundle at the base of the nodule and also the apex of the nodule.

Three germin-like proteins in *A. thaliana* have been characterized to be associated with extracellular matrix (Membré *et al.* 2000). Similarly, rhicadhesin receptor examined in this study may also be associated with extracellular matrix. It was predicted as a secreted protein by bioinformatics tools and is

expressed in vascular bundle as shown by *GUS* reporter gene.

Rhcadhesin receptor from *Pisum sativum*, *PsGER1*, the first known plant germin-like protein to be associated with nodules, is expressed strongly in leaves, stem, roots and nodules, while its expression in flower is weak (Gucciardo *et al.* 2007). The proposed role of rhcadhesin receptor is to mediate the attachment of the bacteria to the plant surface (Swart *et al.* 1994). However, if rhcadhesin receptor (MtRHRE1) examined in this study functions to mediate the attachment of rhizobia to the plant surface, then it should be expressed in the root hairs and epidermis of the whole root, which was not the case. As seen in the Figure 3F, *GUS* expression was detected in the whole nodule and vascular bundle but not in the epidermis of the root, suggesting that MtRHRE1 might not function as attachment protein for rhizobia.

Lipid Transfer Proteins (MtN5 and MtNsLTP). In *P. sativum*, a lipid transfer protein was expressed in the region of expanding cells just proximal to the apical meristem and was expressed weakly in the epidermis (Gucciardo *et al.* 2007). A non-specific lipid transfer protein in *M. truncatula*, MtN5, is proposed to be involved more in nodule organogenesis than in nitrogen fixation for two reasons. First, it is expressed in uninfected Nar nodules (nodulation in the absence of *Rhizobium*) (Truchet *et al.* 1989). Second, its expression in early nodulation before nitrogen fixation takes place is stronger than its expression in mature nodules. In this study, promoter of *MtN5* could not be cloned and only transformant roots expressing *GUS* under *MtNsLTP* promoter were generated. Figure 3C showed that *GUS* was expressed in root vascular bundle and the whole body of 14-day old nodule with stronger expression in the apex zone. It is possible that *MtNsLTP* might have general role in nodule development.

Quantitative real time RT-PCR showed that *MtN5* and *MtNsLTP* (non-specific lipid transfer protein) were both expressed higher in roots and root tips than in aerial organs (Figure 1C & D). Moreover, *MtN5* expression was significantly (P value < 0.05) up-regulated (15-fold) in inoculated root segments relative to control root segment indicating that MtN5 may be involved in nodule development or host defense during nodulation.

Secreted Proteins are Expressed Stronger in Vascular Bundles. Previous studies identified soluble secreted proteins by isolating them from xylem sap and confirmed that they are secreted proteins by identifying their N-terminal sequences of the mature

proteins that do not contain N-terminal signal sequence (Kehr *et al.* 2005; Alvarez *et al.* 2006; Djordjevic *et al.* 2007). In the previous study (Kusumawati *et al.* 2008), proteins were isolated from the medium of cell suspension culture and were identified using MALDI TOF-TOF and searching MtGI database. Using bioinformatic programs, the identified proteins were predicted to be secreted.

The use of promoter:*GUS* constructs, showed that all six promoters examined were strongly expressed in vascular bundles and root tips. They were *MtSBT1*, *MtSCP1*, *MtNsLTP*, *MtFAD-BP1*, *MtRHRE1*, and *MtENOD-like2*. Their strong expression in the vascular bundle suggest that they might be involved in vascular bundle development or function and that the gene products would be released into apoplast on a continuous basis along the entire length of the vasculature. Their strong expression in the vascular bundle might also suggest that these six secreted proteins are secreted to the cells in the vasculature and therefore it is likely that they might become part of the xylem sap. This conclusion is supported by the literature. Subtilisin-like serine protease was found in the xylem sap of *Glycine max* (Subramanian *et al.* 2009), *Zea mays* (Alvarez *et al.* 2006), *Brassica napus*, *Brassica oleracea*, *Cucurbita maxima*, and *Cucumis sativus* (Buhtz *et al.* 2004). Serine carboxypeptidase was among the xylem sap proteins isolated from *Brassica napus* (Kehr *et al.* 2005). A lipid transfer protein was detected in xylem sap of *Glycine max* (Djordjevic *et al.* 2007).

In conclusion, the expression localization study carried out using the native promoter:*GUS* fusions could give an indication concerning which tissues expressed the genes encoding secreted proteins. A reasonable assumption would be that these proteins were then secreted directly into the apoplast surrounding these cells. This could be verified either by generating a secreted-protein-*GFP* fusion or by generating secreted protein specific antibodies and determining the location of the secreted proteins using immunolocalization. Using *GUS* reporter gene assay, this study showed that *MtSBT1*, *MtSCP1*, *MtNsLTP*, *MtFAD-BP1*, *MtRHRE1*, and *MtENOD-like2* strongly expressed in vascular bundles.

Using quantitative real time RT-PCR, six genes encoding secreted proteins examined here showed higher expression in root tips than in leaves, cotyledons or root segments. These six genes were *MtSBT1*, *MtSCP1*, *MtN5*, *MtNsLTP*, *MtFAD-BP1*, and *MtRHRE1*. Only *MtENOD2-like* showed lower expression in roots (either inoculated or non-inoculated) than in leaves and cotyledons (Figure 1).

ACKNOWLEDGEMENT

Lucia Kusumawati was supported by an Australian Development Scholarship. Research was conducted in Research School of Biology, The Australian National University.

REFERENCES

- Alvarez S, Goodger JQD, Marsh EL, Chen S, Asirvatham VS, Schachtman DP. 2006. Characterization of the maize xylem sap proteome. *J Proteome Res* 5:963-972. <http://dx.doi.org/10.1021/pr050471q>
- Bisseling T, Franssen HJ, Govers F, Horvath B, Moerman M, Scheres B, Van de Wiel C, Yang WC. 1990. Early nodulin gene expression during the pea-*Rhizobium* symbiosis. Dordrecht: Kluwer Academic.
- Boisson-Dernier A, Chabaud M, Garcia F, Becard G, Rosenberg C, Barker DG. 2001. *Agrobacterium rhizogenes*-transformed roots of *Medicago truncatula* for the study of nitrogen-fixing and endomycorrhizal symbiotic associations. *Mol Plant-Microbe Interact* 14:695-700. <http://dx.doi.org/10.1094/MPMI.2001.14.6.695>
- Buhtz A, Kolasa A, Arlt K, Walz C, Kehr J. 2004. Xylem sap protein composition is conserved among different plant species. *Planta* 219:610-618. <http://dx.doi.org/10.1007/s00425-004-1259-9>
- Caetano-Anolles G, Gresshoff PM. 1991. Plant genetic control of nodulation. *Annu Rev Microbiol* 45:345-382. <http://dx.doi.org/10.1146/annurev.mi.45.100191.002021>
- Casamitjana-Martinez E, Hofhuis HF, Xu J, Liu CM, Heidstra R, Scheres B. 2003. Root-specific *CLE19* overexpression and the *sol1/2* suppressors implicate a CLV-like pathway in the control of *Arabidopsis* root meristem maintenance. *Curr Biol* 13:1435-1441. [http://dx.doi.org/10.1016/S0960-9822\(03\)00533-5](http://dx.doi.org/10.1016/S0960-9822(03)00533-5)
- Clark SE, Williams RW, Meyerowitz EM. 1997. The CLAVATA1 gene encodes a putative receptor kinase that controls shoot and floral meristem size in *Arabidopsis*. *Cell* 89:575-585. [http://dx.doi.org/10.1016/S0092-8674\(00\)80239-1](http://dx.doi.org/10.1016/S0092-8674(00)80239-1)
- Djordjevic MA, Oakes M, Li DX, Hwang CH, Hocart CH, Gresshoff PM. 2007. The *Glycine max* xylem sap and apoplast proteome. *J Proteome Res* 6:3771-3779. <http://dx.doi.org/10.1021/pr0606833>
- Ehrhardt DW, Atkinson EM, Faull KF, Freedberg DI, Sutherland DP, Armstrong R, Long SR. 1995. In vitro sulfotransferase activity of NodH, a nodulation protein of *Rhizobium meliloti* required for host-specific nodulation. *J Bacteriol* 177:6237-6245.
- Fähraeus G. 1957. The infection of clover root hairs by nodule bacteria studied by a simple glass technique. *J Gen Microbiol* 16:374-381. <http://dx.doi.org/10.1099/00221287-16-2-374>
- Fecht-Christoffers MM, Braun HP, Lemaitre-Guillier C, VanDorselaer A, Horst WJ. 2003. Effect of manganese toxicity on the proteome of the leaf apoplast in cowpea. *Plant Physiol* 133:1935-1946. <http://dx.doi.org/10.1104/pp.103.029215>
- Fletcher JC, Brand U, Running MP, Simon R, Meyerowitz EM. 1999. Signaling of cell fate decisions by CLAVATA3 in *Arabidopsis* shoot meristems. *Science* 283:1911-1914. <http://dx.doi.org/10.1126/science.283.5409.1911>
- Foster CM, Horner HT, Graves WR. 2000. Accumulation of ENOD2-like transcripts in non-nodulating woody Papilionoid legumes. *Plant Physiol* 124:741-750. <http://dx.doi.org/10.1104/pp.124.2.741>
- Franklin-Tong N, Franklin FCH. 2003. Gametophytic self-incompatibility inhibits pollen tube growth using different mechanisms. *Trends Plant Sci* 8:598-605. <http://dx.doi.org/10.1016/j.tplants.2003.10.008>
- Friedrichsen DM, Joazeiro CAP, Li J, Hunter T, Chory J. 2000. Brassinosteroid-insensitive-1 is a ubiquitously expressed leucine-rich repeat receptor serine/threonine kinase. *Plant Physiol* 123:1247-1256. <http://dx.doi.org/10.1104/pp.123.4.1247>
- Fuller F, Kunstner PW, Nguyen T, Verma DPS. 1983. Soybean nodulin genes: Analysis of cDNA clones reveals several major tissue-specific sequences in nitrogen-fixing root nodules. *Proc Nat Acad Sci USA* 80: 2594-2598. <http://dx.doi.org/10.1073/pnas.80.9.2594>
- Gage DJ, Margolin W. 2000. Hanging by a thread: Invasion of legume plants by rhizobia. *Curr Opin Microbiol* 3:613-617. [http://dx.doi.org/10.1016/S1369-5274\(00\)00149-1](http://dx.doi.org/10.1016/S1369-5274(00)00149-1)
- Gloude-mans T, Bisseling T. 1989. Plant gene expression in early stages of *Rhizobium*-legume symbiosis. *Plant Sci* 65:1-14. [http://dx.doi.org/10.1016/0168-9452\(89\)90201-X](http://dx.doi.org/10.1016/0168-9452(89)90201-X)
- Govers F, Moerman M, Van Kammen A, Bisseling T. 1985. Expression of plant genes during the development of pea root nodules. *EMBO J* 4:861-867.
- Greenbaum D, Luscombe NM, Jansen R, Qian J, Gerstein M. 2001. Interrelating different types of genomic data, from proteome to secretome: Oming in on function. *Genome Res* 11:1463-1468. <http://dx.doi.org/10.1101/gr.207401>
- Grunwald I, Rupprecht I, Schuster G, Kloppstech K. 2003. Identification of guttation fluid proteins: The presence of pathogenesis-related proteins in non-infected barley plants. *Physiol Plant* 119:192-202. <http://dx.doi.org/10.1034/j.1399-3054.2003.00202.x>
- Gualtieri G, Bisseling T. 2000. The evolution of nodulation. *Plant Mol Biol* 42:181-194. <http://dx.doi.org/10.1023/A:1006396525292>
- Gucciardo S, Wisniewski JP, Brewin NJ, Bornemann S. 2007. A germin-like protein with superoxide dismutase activity in pea nodules with high protein sequence identity to a putative rhicadhesin receptor. *J Ex Bot* 58:1161-1171. <http://dx.doi.org/10.1093/jxb/erl282>
- Gulash M, Ames P, Larosiliere RC, Bergman K. 1984. Rhizobia are attracted to localized sites on legume roots. *Appl Environ Microbiol* 48:149-152.
- He Z, Wang ZY, Li J, Zhu Q, Lamb C, Ronald P, Chory J. 2000. Perception of brassinosteroids by the extracellular domain of the receptor kinase BRI1. *Science* 288:2360-2363. <http://dx.doi.org/10.1126/science.288.5475.2360>
- Höfgen R, Willmitzer L. 1988. Storage of competent cells for *Agrobacterium* transformation. *Nucleic Acids Res* 16:9877. <http://dx.doi.org/10.1093/nar/16.20.9877>
- Holmes P, Goffard N, Weiller G, Rolfe B, Imin N. 2008. Transcriptional profiling of *Medicago truncatula* meristematic root cells. *BMC Plant Biol* 8:21. <http://dx.doi.org/10.1186/1471-2229-8-21>
- Hugot K, Riviere MP, Moreillon C, Dayem MA, Cozzitorto J, Arbiol G, Barbry P, Weiss C, Galiana E. 2004. Coordinated regulation of genes for secretion in tobacco at late developmental stages: Association with resistance against oomycetes. *Plant Physiol* 134:858-870. <http://dx.doi.org/10.1104/pp.103.034173>

- Jeong S, Trotochaud AE, Clark SE. 1999. The *Arabidopsis* CLAVATA2 gene encodes a receptor-like protein required for the stability of the CLAVATA1 receptor-like kinase. *Plant Cell* 11:1925-1934. <http://dx.doi.org/10.1105/tpc.11.10.1925>
- Jones RL, Robinson DG. 1989. Tansley Review No. 17. Protein secretion in plants. *New Phytol* 111:567-597. <http://dx.doi.org/10.1111/j.1469-8137.1989.tb02352.x>
- Kape R, Parniske M, Werner D. 1991. Chemotaxis and nod gene activity of *Bradyrhizobium japonicum* in response to hydroxycinnamic acids and isoflavonoids. *Appl Environ Microbiol* 57:316-319.
- Kehr J, Buhtz A, Giavalisco P. 2005. Analysis of xylem sap proteins from *Brassica napus*. *BMC Plant Biol* 5:11. <http://dx.doi.org/10.1186/1471-2229-5-11>
- Krusell L, Madsen LH, Sato S, Aubert G, Genua A, Szczyglowski K, Duc G, Kaneko T, Tabata S, de Bruijn F, Pajuelo E, Sandal N, Stougaard J. 2002. Shoot control of root development and nodulation is mediated by a receptor-like kinase. *Nature* 420:422-426. <http://dx.doi.org/10.1038/nature01207>
- Kusumawati L, Imin N, Djordjevic MA. 2008. Characterization of the secretome of suspension cultures of *Medicago* species reveals proteins important for defense and development. *J Proteome Res* 7:4508-4520. <http://dx.doi.org/10.1021/pr800291z>
- Lang-Unnasch N, Ausubel FM. 1985. Nodule-specific polypeptides from effective alfalfa root nodules and from ineffective nodules lacking nitrogenase. *Plant Physiol* 77:833-839. <http://dx.doi.org/10.1104/pp.77.4.833>
- Laplaze L, Ribeiro A, Franche C, Duhoux E, Auguy F, Bogusz D, Pawlowski K. 2000. Characterization of a *Casuarina glauca* nodule-specific subtilisin-like protease gene, a homolog of *Alnus glutinosa* ag12. *Mol Plant-Microbe Interact* 13:113-117. <http://dx.doi.org/10.1094/MPMI.2000.13.1.113>
- Lee SJ, Saravanan RS, Damasceno CMB, Yamane H, Kim BD, Rose JKC. 2004. Digging deeper into the plant cell wall proteome. *Plant Physiol Bioch* 42:979-988. <http://dx.doi.org/10.1016/j.plaphy.2004.10.014>
- Li J, Chory J. 1997. A putative leucine-rich repeat receptor kinase involved in brassinosteroid signal transduction. *Cell* 9:929-938. [http://dx.doi.org/10.1016/S0092-8674\(00\)80357-8](http://dx.doi.org/10.1016/S0092-8674(00)80357-8)
- Li J, Lease KA, Tax FE, Walker JC. 2001. BRS1, a serine carboxypeptidase, regulates BRI1 signaling in *Arabidopsis thaliana*. *Proc Nat Acad Sci USA* 98:5916-5921. <http://dx.doi.org/10.1073/pnas.091065998>
- Lorquin J, Lortet G, Ferro M, Méar N, Dreyfus B, Promé JC, Boivin C. 1997. Nod factors from *Sinorhizobium saheli* and *S. teranga* bv. *sesbaniae* are both arabinosylated and fucosylated, a structural feature specific to *Sesbania rostrata* symbionts. *Mol Plant-Microbe Interact* 10:879-890. <http://dx.doi.org/10.1094/MPMI.1997.10.7.879>
- Martínez-Abarca F, Herrera-Cervera JA, Bueno P, Sanjuan J, Bisseling T, Olivares J. 1998. Involvement of salicylic acid in the establishment of the *Rhizobium meliloti*-alfalfa symbiosis. *Mol Plant-Microbe Interact* 11:153-155. <http://dx.doi.org/10.1094/MPMI.1998.11.2.153>
- Matsubayashi Y, Sakagami Y. 1996. Phytosulfokine, sulfated peptides that induce the proliferation of single mesophyll cells of *Asparagus officinalis* L. *Proc Nat Acad Sci USA* 93:7623-7627. <http://dx.doi.org/10.1073/pnas.93.15.7623>
- Matsubayashi Y, Takagi L, Sakagami Y. 1997. Phytosulfokine- α , a sulfated pentapeptide, stimulates the proliferation of rice cells by means of specific high- and low-affinity binding sites. *Proc Nat Acad Sci USA* 94:13357-13362. <http://dx.doi.org/10.1073/pnas.94.24.13357>
- Matsubayashi Y, Takagi L, Omura N, Morita A, Sakagami Y. 1999. The endogenous sulfated pentapeptide phytosulfokine- α stimulates tracheary element differentiation of isolated mesophyll cells of *Zinnia*. *Plant Physiol* 120:1043-1048. <http://dx.doi.org/10.1104/pp.120.4.1043>
- Membré N, Bernier F, Staiger D, Berna A. 2000. *Arabidopsis thaliana* germin-like proteins: common and specific features point to a variety of functions. *Planta* 211:345-354. <http://dx.doi.org/10.1007/s004250000277>
- Ndimba BK, Chivasa S, Hamilton JM, Simon WJ, Slabas AR. 2003. Proteomic analysis of changes in the extracellular matrix of *Arabidopsis* cell suspension cultures induced by fungal elicitors. *Proteomics* 3:1047-1059. <http://dx.doi.org/10.1002/pmic.200300413>
- Niehaus K, Kapp D, Pühler A. 1993. Plant defence and delayed infection of alfalfa pseudonodules induced by an exopolysaccharide (EPS I)-deficient *Rhizobium meliloti* mutant. *Planta* 190:415-425. <http://dx.doi.org/10.1007/BF00196971>
- Nishimura R, Hayashi M, Wu GJ, Kouchi H, Imaizumi-Anraku H, Murakami Y, Kawasaki S, Akao S, Ohmori M, Nagasawa M, Harada K, Kawaguchi M. 2002. HAR1 mediates systemic regulation of symbiotic organ development. *Nature* 420:426-429. <http://dx.doi.org/10.1038/nature01231>
- Noguchi T, Fujioka S, Choe S, Takatsuto S, Yoshida S, Yuan H, Feldmann KA, Tax FE. 1999. Brassinosteroid-Insensitive dwarf mutants of *Arabidopsis* accumulate brassinosteroids. *Plant Physiol* 121:743-752. <http://dx.doi.org/10.1104/pp.121.3.743>
- Nukui N, Ezura H, Yuhashi KI, Yasuta T, Minamisawa K. 2000. Effects of ethylene precursor and inhibitors for ethylene biosynthesis and perception on nodulation in *Lotus japonicus* and *Macroptilium atropurpureum*. *Plant Cell Physiol* 41:893-897. <http://dx.doi.org/10.1093/pcp/pcd011>
- Oldroyd GED, Downie JA. 2008. Coordinating nodule morphogenesis with rhizobial infection in legumes. *Ann Rev Plant Biol* 59:519-546. <http://dx.doi.org/10.1146/annurev.arplant.59.032607.092839>
- Pearce G, Moura DS, Stratmann J, Ryan CA. 2001. RALF, a 5-kDa ubiquitous polypeptide in plants, arrests root growth and development. *Proc Nat Acad Sci USA* 98:12843-12847. <http://dx.doi.org/10.1073/pnas.201416998>
- Perotto S, Brewin NJ, Kannenberg EL. 1994. Cytological evidence for a host defense response that reduces cell and tissue invasion in pea nodules by lipopolysaccharide-defective mutants of *Rhizobium leguminosarum* Strain 3841. *Mol Plant-Microbe Interact* 7:99-112. <http://dx.doi.org/10.1094/MPMI-7-0099>
- Peters NK, Frost JW, Sharon RL. 1986. A plant flavone, luteolin, induces expression of *Rhizobium meliloti* nodulation genes. *Science* 233:977-980. <http://dx.doi.org/10.1126/science.3738520>
- Prayitno J, Imin N, Rolfe BG, Mathesius U. 2006b. Identification of ethylene-mediated protein changes during nodulation in *Medicago truncatula* using proteome analysis. *J Proteome Res* 5:3084-3095. <http://dx.doi.org/10.1021/pr0602646>

- Prayitno J, Rolfe BG, Mathesius U. 2006a. The Ethylene-Insensitive sickle Mutant of *Medicago truncatula* Shows Altered Auxin Transport Regulation during Nodulation. *Plant Physiol* 142:168-180. <http://dx.doi.org/10.1104/pp.106.080093>
- Redmond JW, Batley M, Djordjevic MA, Innes RW, Kuempel PL, Rolfe BG. 1986. Flavones induce expression of nodulation genes in *Rhizobium*. *Nature* 323:632-635. <http://dx.doi.org/10.1038/323632a0>
- Rep M, Dekker HL, Vossen JH, de Boer AD, Houterman PM, Speijer D, Back JW, de Koster CG, Cornelissen BJC. 2002. Mass spectrometric identification of isoforms of PR proteins in xylem sap of fungus-infected tomato. *Plant Physiol* 130:904-917. <http://dx.doi.org/10.1104/pp.007427>
- Ribeiro A, Akkermans ADL, van Kammen A, Bisseling T, Pawlowski K. 1995. A nodule-specific gene encoding a subtilisin-like protease is expressed in early stages of actinorhizal nodule development. *Plant Cell* 7:785-794. <http://dx.doi.org/10.1105/tpc.7.6.785>
- Roche P, Debellé F, Maillat F, Lerouge P, Faucher C, Truchet G, Dénarié J, Promé JC. 1991. Molecular basis of symbiotic host specificity in *Rhizobium meliloti*: nodH and nodPQ genes encode the sulfation of lipo-oligosaccharide signals. *Cell* 67:1131-1143. [http://dx.doi.org/10.1016/0092-8674\(91\)90290-F](http://dx.doi.org/10.1016/0092-8674(91)90290-F)
- Rolfe BG, Gresshoff PM, Shine J. 1980. Rapid screening for symbiotic mutants of *Rhizobium* and white clover. *Plant Sci Lett* 19:277-284. [http://dx.doi.org/10.1016/0304-4211\(80\)90082-6](http://dx.doi.org/10.1016/0304-4211(80)90082-6)
- Sanderfoot AA, Raikhel NV. 2003. The secretory system of *Arabidopsis*. In: Somerville CR, Meyerowitz EM (eds). *The Arabidopsis Book*. Rockville: American Society of Plant Biologists.
- Scheres B, McKhann HI, Zalensky A, Lobler M, Bisseling T, Hirsch AM. 1992. The *PsENOD12* gene is expressed at two different sites in Afghanistan pea pseudonodules induced by auxin transport inhibitors. *Plant Physiol* 100:1649-1655. <http://dx.doi.org/10.1104/pp.100.4.1649>
- Schnabel E, Etienne-Pascal J, de Carvalho-Niebel F, Duc G, Frugoli J. 2005. The *Medicago truncatula* *SUNN* gene encodes a CLV1-like leucine-rich repeat receptor kinase that regulates nodule number and root length. *Plant Mol Biol* 58:809-822. <http://dx.doi.org/10.1007/s11103-005-8102-y>
- Searle IR, Men AE, Laniya TS, Buzas DM, Iturbe-Ormaetxe I, Carroll BJ, Gresshoff PM. 2003. Long-distance signaling in nodulation directed by a CLAVATA1-like receptor kinase. *Science* 299:109-112. <http://dx.doi.org/10.1126/science.1077937>
- Streeter J, Wong PP. 1988. Inhibition of legume nodule formation and N₂ fixation by nitrate. *CRC Crit Rev Plant Sci* 7:1-23. <http://dx.doi.org/10.1080/07352688809382257>
- Subramanian S, Cho UH, Keyes C, Yu O. 2009. Distinct changes in soybean xylem sap proteome in response to pathogenic and symbiotic microbe interactions. *BMC Plant Biol* 9:119. <http://dx.doi.org/10.1186/1471-2229-9-119>
- Swart S, Logman TJJ, Smit G, Lugtenberg BJJ, Kijne JW. 1994. Purification and partial characterization of a glycoprotein from pea (*Pisum sativum*) with receptor activity for rhicadhesin, an attachment protein of *Rhizobiaceae*. *Plant Mol Biol* 24:171-183. <http://dx.doi.org/10.1007/BF00040583>
- Szczyglowski K, Hamburger D, Kapranov P, de Bruijn FJ. 1997. Construction of a *Lotus japonicus* late nodulin expressed sequence tag library and identification of novel nodule-specific genes. *Plant Physiol* 114:1335-1346. <http://dx.doi.org/10.1104/pp.114.4.1335>
- Truchet G, Barker DG, Camut S, Billy F, Vasse J, Huguet T. 1989. Alfalfa nodulation in the absence of *Rhizobium*. *Mol Gen Genet* 219:65-68. <http://dx.doi.org/10.1007/BF00261158>
- van de Wiel C, Scheres B, Franssen H, van Lierop MJ, van Lammeren A, van Kammen A, Bisseling T. 1990. The early nodulin transcript ENOD2 is located in the nodule parenchyma (inner cortex) of pea and soybean root nodules. *EMBO J* 9:1-7.
- van Rhijn P, Fang Y, Galili S, Shaul O, Atzmon N, Winger S, Eshed Y, Lum M, Li Y, To V, Fujishige N, Kapulnik Y, Hirsch AM. 1997. Expression of early nodulin genes in alfalfa mycorrhizae indicates that signal transduction pathways used in forming arbuscular mycorrhizae and *Rhizobium*-induced nodules may be conserved. *Proc Nat Acad Sci USA* 94:5467-5472. <http://dx.doi.org/10.1073/pnas.94.10.5467>
- Vitha S, Beneš K, Phillips JP, Gartland KMA. 1995. Histochemical GUS Analysis. In: Gartland KMA, Davey MR (eds). *Agrobacterium Protocols*. Totowa, New Jersey: Humana Pr. p 185-193. <http://dx.doi.org/10.1385/0-89603-302-3:185>
- von Groll U, Berger D, Altmann T. 2002. The subtilisin-like serine protease SDD1 mediates cell-to-cell signaling during *Arabidopsis* stomatal development. *Plant Cell* 14:1527-1539. <http://dx.doi.org/10.1105/tpc.001016>
- Wasson AP, Pellerone FI, Mathesius U. 2006. Silencing the flavonoid pathway in *Medicago truncatula* inhibits root nodule formation and prevents auxin transport regulation by rhizobia. *Plant Cell* 18:1617-1629. <http://dx.doi.org/10.1105/tpc.105.038232>
- Wopereis J, Pajuelo E, Dazzo FB, Jiang Q, Gresshoff PM, Bruijn FJD, Stougaard J, Szczyglowski K. 2000. Short root mutant of *Lotus japonicus* with a dramatically altered symbiotic phenotype. *Plant J* 23:97-114. <http://dx.doi.org/10.1046/j.1365-313x.2000.00799.x>
- Zhou A, Li J. 2005. *Arabidopsis* BRS1 is a secreted and active serine carboxypeptidase. *J Biol Chem* 280:35554-35561. <http://dx.doi.org/10.1074/jbc.M503299200>