

The Growth of Giant Clams Juvenil is Influenced by Nutrient Addition

Pengaruh Penambahan Nutrien terhadap Pertumbuhan Juvenil Kima

Ambariyanto

Marine Science Department, FPIK, Diponegoro University
Kampus Tembalang, Semarang. E-mail: ambariyanto@telkom.net

Abstrak

Ekosistem karang merupakan daerah yang memiliki produktivitas tinggi namun kandungan nutrient anorganiknya sangat rendah. Sehingga dapat diduga bahwa kenaikan nutrient anorganik dalam ekosistem ini akan mempengaruhi kehidupan organisme yang ada di dalamnya. Penelitian ini bertujuan untuk melihat pengaruh peningkatan nutrien yakni N dan P terhadap pertumbuhan juvenil Kima. Hasil penelitian menunjukkan bahwa dengan penambahan nutrien N secara nyata meningkatkan pertumbuhan juvenil Kima tersebut. Di lain pihak penambahan P tidak berpengaruh terhadap laju pertumbuhan juvenil Kima. Hasil ini menunjukkan bahwa penambahan ammonium seharusnya dilakukan di hatchery untuk meningkatkan pertumbuhan dan produksi Kima.

Kata Kunci: Kima, *Tridacna squamosa*, nutrien, ammonium, fosfat

Diterima: 13 Januari 2007, disetujui: 09 April 2007

Introduction

There is an important fact that coral reef ecosystem is known to have high productivity, although the concentrations of inorganic nutrients within the waters are very low. The nutrient recycling between zooxanthellae (single cell dinoflagellate algae live in association with many marine invertebrates) and their hosts is thought to be the one which responsible this contradictive condition.

Zooxanthellae and their hosts (such as corals, anemone, giant clams, etc) are known to be able to absorb nutrients from their surrounding waters (Miller & Yellowlees, 1989). The existence of absorption capability by zooxanthellae and their host and the fact that coral reef ecosystems are known to have low concentration of nutrients, expectation that increasing nutrient concentrations in the water should have a significant impact on the invertebrate-zooxanthellae symbiosis. Increasing the concentration of inorganic nutrients leads to changes in the biochemistry,

physiology and morphology of symbiotic invertebrates (Dubinsky *et al.*, 1990; Fitt *et al.*, 1993; Hoegh-Guldberg, 1994).

The changes due to increasing nutrient concentration occurred in zooxanthellae alone or in both the zooxanthellae and the host (Hoegh-Guldberg & Smith, 1989; Hastie *et al.*, 1992; Belda *et al.*, 1993a,b; Achituv *et al.*, 1994; Muller-Parker *et al.*, 1994a,b). One of the important effects of high nutrient concentration worth mention is increasing number of zooxanthellae within the host (Ambariyanto, 1999).

Since the growth of giant clams, known as a member of coral reef community, is influenced by zooxanthellae through photosynthesis translocation, therefore it suggests that increasing nutrient concentration would improve giant clams growth. The present study aims to investigate the effects of nutrient addition on the growth of juvenile of giant clams (*Tridacna squamosa*).

Material and Methods

Twelve glass aquariums (1 L) were used in the present study. These aquariums were thoroughly cleaned using mild concentration of chlorin and were rinsed with clean fresh and seawater for at least ten times before being used for the experiment. These aquariums then were filled with filtered seawater (using filter bag) and also 10 of three months old giant clams juvenil with approximately 5 mm shell length. Aeration was given during the experiment in order to alleviate dissolved oxygen content within the water.

Three treatments (inorganic nutrient addition i.e. N, P and N+P) and control (without any nutrient addition) were carried out with three replicates. Final concentration of nutrient given were 20 μ M for N (ammonium; NH₄Cl) and 2 μ M for P (phosphorus; KH₂PO₄). Giant clams shell length were measured using callipers to the nearest of 0.1 mm before and after the experiment. This experiment was carried out for seven days.

Analysis of variance (ANOVA) was used to analyse data of percentage giant clam growth (shell length). Normality and homogeneity of variances was tested prior to ANOVA. Differences among the means were

tested using the Student-Newman-Keuls test (Underwood, 1981).

Results and Discussions

The results of the present study show that addition of inorganic nutrient, especially N (ammonium) increased the shell length of giant clams juvenil. Compared with control, the shell length increment of N treated clams were almost seven times higher. The influenced of the addition of P (phosphorus), however, was much less than that of N addition (almost 9 fold less).

Inorganic nutrient addition, especially N, increase the growth rate of giant clams juvenil. Figure 1. shows juvenile which grown in the water added with N have significantly longer shell length compared with those other treatments and control (P= 4.6E-06).

Coral reef ecosystem, where giant clam populations can be found has low concentration of inorganic nutrients, although this ecosystem is known to be very productive. Therefore, changes in the concentration of these nutrients would change the biological processes of the organisms which live in the ecosystem.




Figure 1. Mean \pm SD percentage of giant clams growth (shell length) treated with different nutrient addition

Giant clams biological processes are influenced by the concentration of anorganic nutrients N (ammonium) and P (phosphate) in the water. Scientists have found that higher the growth rate of zooxanthellae within the mantle of the clams was found on the clams added by nutrients especially N. On the other hand, the concentration of P did not significantly influenced zooxanthellae growth rate (Ambariyanto & Hoegh-Guldberg, 1997), but affects the structure of giant clams shells (Belda *et al.*, 1993b). Several authors have found that increasing the concentrations of inorganic nutrients does not influence coral biomass characteristics, but changes a number of aspects of zooxanthellae biomass, such as population density, mitotic index, chlorophyll *a* content, and C:N ratio of zooxanthellae (Hoegh-Guldberg & Smith, 1989; Muscatine *et al.*, 1989; Stambler *et al.*, 1991).

The results of the present study show that higher concentration of nutrient (N) increased giant clam juvenil growth rate ($P=4.6E-06$). See Figure 1. This result is in accordance with other reports which showed that nutrient addition in the water have increased the number of corals zooxanthellae (Hoegh-Guldberg 1994), as well as adult giant clams, *Tridacna maxima* (Ambariyanto & Hoegh-Guldberg, 1997).

Other reports showed that giant clam, *Tridacna gigas* responds to inorganic nutrient enrichment through an increase in zooxanthellae density, chlorophyll *a* content, and also soft tissue weight, C:N and C:P ratio of the soft tissue, and the growth rate of the clams (Braley *et al.*, 1992; Belda *et al.*, 1993b; Fitt *et al.*, 1993). Similar results were found in coral, *Pocillopora damicornis* exposed to elevated ammonium concentrations showed increased in zooxanthellae density, chlorophyll *a* content and protein content of the coral (Muller-Parker *et al.*, 1994b). However, in the same experiment, Achituv *et al.*, (1994) found no changes in the biochemical composition of the coral tissue due to ammonium enrichment, but it changed the lipid and protein content of zooxanthellae.

Ambariyanto (1996) found that, nutrient enrichment resulted in an increase in

zooxanthellae density and chlorophyll *a* concentration, and by changes in the ultrastructure of zooxanthellae. No changes were observed on the animal biomass parameters measured (*ie.* soft tissue weight, C:N ratio and protein content of the mantle of the clam). A similar phenomenon has been reported by Muller-Parker *et al.*, (1994b) who found changes in C:N ratio of zooxanthellae but not in the tissue of the coral *Pocillopora damicornis* due to ammonium enrichment.

Increasing the number of zooxanthellae has resulted in the increment of giant clams growth rate due to higher amount of photosynthates being translocated into the host. The results of the present study suggests that nutrient addition into the medium (seawater) where the juvenile are grown should be done in the hatchery before outgrow in the ocean.

Conclusion

Increasing inorganic nutrient especially N (ammonium) significantly improved the growth (in shell length) giant clams juvenil. It is suggested that in daily operational of giant clams hatchery, nutrient enrichment should not only be done during larval development (until metamorphoses), but also at juvenil stage.

Acknowledgements

This study was funded by RUT (Riset Unggulan Terpadu). The author would like to thank the management of Barang Lompo Island Marine Lab. UNHAS, Makassar. This work would not be possible without any help from Dr. Iqbal Jawad, Dr. Niartiningsih, Ihsan Ramli ST, Pak Ipul, Pudir, Japri, and Pak Ridwan and his family.

References

- Achituv, Y., Ben-Zion, M. and Mizrahi, L. 1994. Carbohydrate, lipid, and protein composition of zooxanthellae and animal fractions of the coral *Pocillopora damicornis* exposed to ammonium enrichment. *Pac. Sci.* 48 (3): 224-233.

The Growth of Giant Clams Juvenil

- Ambariyanto. 1996. Effects of nutrient enrichment in the field on the giant clam, *Tridacna maxima*. PhD Thesis. The University of Sydney, Sydney Australia. 267 p.
- Ambariyanto and Hoegh-Guldberg, O. 1997. The effects of nutrient enrichment on the biomass, growth and calcification of giant clam, *Tridacna maxima*. *Mar. Biol.* 129 (4): 635-642.
- Ambariyanto. 1999. Enrichment of inorganic nutrients: Biological consequences of giant clams (Tridacnidae). *J. Coast. Dev.* 2 (3): 435-441
- Belda, C.A., Lucas, J.S. and Yellowlees, D. 1993a. Nutrient limitation in the giant clam-zooxanthellae symbiosis: effects of nutrient supplements on growth of the symbiotic partners. *Mar. Biol.* 117: 655-664.
- Belda, C.A., Cuff, C. and Yellowlees, D. 1993b. Modification of shell formation in the giant clam *Tridacna gigas* at elevated nutrient levels in sea water. *Mar. Biol.* 117: 251-257.
- Braley, R.D., Sutton, D., Mingo, S.S.M. and Southgate, P.C. 1992. Passive greenhouse heating, recirculation, and nutrient addition for nursery phase *Tridacna gigas*: growth boost during winter months. *Aquaculture* 108: 29-50.
- Dubinsky, Z., Stambler, N., Ben-Zion, M. and McCloskey, L.R. 1990. The effect of external resources of the optical properties and photosynthetic efficiency of *Stylophora pistillata*. *Proc. R. Soc. (Lond) Ser. B.* 239: 231-246
- Fitt, W.K., Heslinga, G.A. and Watson, T.C. 1993. Utilization of dissolved inorganic nutrients in growth and mariculture of the tridacnid clam *Tridacna derasa*. *Aquaculture* 109: 27-38
- Hastie, L.C., Watson, T.C., Isamu, T. and Heslinga, G.A. 1992. Effect of nutrient on *Tridacna derasa* seed: dissolved inorganic nitrogen increases growth rate. *Aquaculture* 106: 41-49.
- Hoegh-Guldberg, O. 1994. Population dynamics of symbiotic zooxanthellae in the coral *Pocillopora damicornis* exposed to elevated ammonium [(NH₄)₂SO₄] concentrations. *Pac. Sci.* 48 (3): 263-272.
- Hoegh-Guldberg, O. and Smith, G.J. 1989. Influence of the population density of zooxanthellae and supply of ammonium on the biomass and metabolic characteristics of the reef corals *Seriatopora hystrix* and *Stylophora pistillata*. *Mar. Ecol. Prog. Ser.* 57: 173-186
- Miller, D.J. and Yellowlees, D. 1989. Inorganic nitrogen uptake by symbiotic marine cnidarians: a critical review. *Proc. R. Soc. Lond. B* 237: 109-125.
- Muller-Parker, G., Cook, C.B. and D'Elia, C.F. 1994a. Elemental composition of the coral *Pocillopora damicornis* exposed to elevated seawater ammonium. *Pac. Sci.* 48 (3): 234-246.
- Muller-Parker, G., McCloskey, L.R., Hoegh-Guldberg, O. and McAuley, P.J. 1994b. Effect of ammonium enrichment on animal and algal biomass of the coral *Pocillopora damicornis*. *Pac. Sci.* 48 (3): 273-283.
- Muscantine, L., Falkowski, P.G., Dubinsky, Z., Cook, P.A. and McCloskey, L.R. 1989. The effect of external nutrient resources on the population dynamics of zooxanthellae in a reef coral. *Proc. R. Soc. Lond. B* 236: 311-324.
- Stambler, N., Popper, N., Dubinsky, Z. and Stimson, J. 1991. Effect of nutrient enrichment and water motion on the coral *Pocillopora damicornis*. *Pac. Sci.* 45 (3): 299-307.
- Underwood, A.J. 1981. Techniques of analysis of variance in experimental marine biology and ecology. *Oceanogr. Mar. Biol. Ann. Rev.* 19: 513-605.