

Meristematic shoot tip culture obtain good quality seedling of citrus (*Citrus nobilis* var. brastepu) free from citrus vein phloem degeneration

Isnaini Nurwahyuni¹, Justin A Napitupulu¹, Rosmayati¹, Fauziah Harahap²

¹Doctorate Programme, Department of Agriculture, Faculty of Agriculture, North Sumatra University, Medan, North Sumatera, Indonesia 20155 E-mail: isnaininurwahyuni@yahoo.co.id. ²Departement Biologi, FMIPA, State University of Medan, Jl. Willem Iskandar, Psr V, Medan, Sumatera Utara, Indonesia 20221.

Abstract. Meristematic shoot tip culture to obtain good quality seedling of citrus (*Citrus nobilis* Var. Brastepu) free from citrus vein phloem degeneration (CVPD) is explained in this research. The method is conducted by using a diseased plant and is then propagated in optimum conditions. The research is conducted in Tissue Culture Laboratory Department of Biology, University of North Sumatra, Indonesia. Optimum conditions for citrus propagation have been obtained to produce good quality of *Citrus* seedlings that is free from CVPD. The analysis of the DNA using PCR method has confirmed that the seedlings are healthy and free from CVPD. Genomic analysis using 10 RAPD markers show the plantlets regenerated from the cultures are 100% similar with the mother plant.

Keywords: Optimization, in vitro propagation, citrus, CVPD free, seedling

Introduction

The propagation of citrus Brastepu are commonly conducted by generative and vegetative methods (Sanford, 1992; Rieger, 2006; Nurwahyuni, *et.al*, 2012). The production of citrus seedling are mostly with vegetative method such as cutting and grafting, (Williamson dan Jackson, 1994; Prastowo, *et.al.*, 2006). However the problem to obtain good root are often found and results in low performance of seedling. The oculation method by joining the bud with lemonade citrus as a mother plant has successfully conducted (Nurwahyuni, *et.al*, 2012). However, the production of the good quality of seedling free from CVPD for mass production is difficult to conduct. Therefore, it is good to find another method to obtain good seedling for mass production by using in vitro propagation. The application of in vitro propagation technique is compulsory to be done as a strategy to conserve local variety citrus in North Sumatra. The genetic potential from local citrus have to be preserved in the quality development of the citrus, such as resistance to plant diseases, quantity and quality of fruits and the content of medicinal bioactive compounds in the plants (Nurwahyuni, 2011). Therefore, the local variety of citrus in North Sumatra has to be conserved. The aim of the research is to apply meristematic shoot tip culture to obtain good quality seedling of citrus free from CVPD. The success in the production of good quality seedling of citrus that is free from CVPD could save the variety local citrus in North Sumatra and will motivate the farmer to grow the citrus in Karo regency, North Sumatera, Indonesia.

In vitro technique has been applied for the propagation of many *dicotyl* plants, and has been conducted for many years to provide seedling of the plants with high economic value (Chaturvedi, *et.all.*, 1974). Various works have been conducted on the use of plant tissue culture to propagate new seedling (Niedz and Evens, 2008) by using various explants such as roots (Grosser, *et.all.*, 2000), leaves, tuber (Costa, *et.all.*, 2004), ovary (Carimi, *et.all.*, 1995), stem and *epicotyl* (Edriss and Burger, 1984) and the protoplast (Das, *et.all.*, 2000). The propagation of *Citrus sinensis* L. Osbeck have been reported (Duran-Vila, *et.all.*, 1989). The effect of various type of growth stimulator into the growth of the callus have also been explained (Maggon and Singh, 1995)). Some factors have to be considered in the tissue culture techniques to obtain good seedling such as the genetic of the plant, the enzymatic reaction, and the metabolism process (Terol, *et.all.*, 2007).

Various variety of local sweet orange planted and developed in Brastagi North Sumatra have been identified, they are Brastepu, Boci, and Rimokeling. The variety Brastepu is known as "Jeruk Brastagi". The variety Brastepu has superior in the genetic as it has very sweet taste compare with honey citrus, the color of a raped feel is reddish, and the fruit is large size fruit. The survey has shown that Citrus Brastepu is very popular in North Sumatra because

the citrus could be used as sources of fruit and the leaves and the feels contains bioactive that is used as medicine to cure some diseases. The seedling for Brastepu is provided by cut and budding technique to obtain the good quality seedling. However, propagation by using cut and budding techniques only produced very limited seedling and therefore it is difficult to provide lage number of seedling for plantation. To overcome the problem, in vitro propagation is the best choice for mass production of good quality seedling at a short time. The propagation of the local citrus has been conducted by using of stem and young leaves as explants followed by micropropagation (Nurwahyuni, 2003). Variation of growth stimulator in culture medium were very effective to initiate callus. Culture medium containing of 2,4-D was found more effective to initiate the callus compare with other growth stimulator such as NAA, ZI and KI or the combination of IAA, BA and KI. It was found that the MS medium enriched with 2,4-D and KI are able to develop the callus to a plantlet. The seedling obtained in the medium culture has been tested for its tolerance into salinity (Nurwahyuni, 2003), their resistance to fungus, and their ability to adapt with extreme season. All of those studies were conducted to obtain a good quality seedling of *Citrus nobilis* Var. Brastepu.

Materials and Methods

Research methods are consisted of identification and characterization of *Citrus nobilis* Var. Brastepu, and experimental procedures are explained in details (Nurwahyuni, et.all., 2012). Experimental procedures are consisted of preparation citrus explant, preparation of culture medium, sterilization and plantation of the explants, and shoot tip culture of *Citrus nobilis* Var. Brastepu. Diseased plants that still grow well, are chosen as the sources of explants, that were obtained from Bukit Village, Kabupaten Karo, Province North Sumatra. In vitro propagation is set up experimentally with 10 replications by using combination of various variable of D₀ is 2,4 dichlorophenoxyacetic acid or 2,4-D 1,0 mg/l; B₀ is Benzyl Amino Purine or BAP 0,0 mg/l; B₁ is BAP 0,5 mg/l; B₂ is BAP 1,0 mg/l; B₃ is BAP 3,0 mg/l. The pH of the medium is adjusted to pH 5.8 – 6.0, and it is then sterilized at 121 °C, 15 lb for 20 minutes. The meristem used as plant material for the culture shown in Figure 1.




Figure 1. The microscopy slides of the meristem shoot tips of *Citrus nobilis* Var. Brastepu as explants: a,b,c,d are a series of cuttings.

Results and Discussion

Description of *Citrus nobilis* Var. Brastepu

The local citrus that has been studied is *Citrus nobilis* Var. Brastepu. It was collected from farmers in many villages, and from The Department of Agriculture in Kabupaten Karo. All informations about the orange are obtained from the farmer based on their experiences on planting and carrying the citrus. The citrus Brastepu are rarely found in the field because the farmer did not interested in planting it for some reason. The performance of *Citrus* Brastepu and appearance of its fruit is presented in Figure 2.

Callus induction of *Citrus nobilis* Var. Brastepu

It was known that the ability of plant tissue to produce callus was effected by the composition and the concentration of culture medium, the presence of growth stimulator, and the light intensity (Nurwahyuni and Tjondronegoro, 1994). Therefore, the basal MS medium is used to initiate callus in this study. The basal MS consisted of growth stimulators are use as reported in the previous study. It was observed that the callus was grown in the culture where the texture of the callus in the medium is almost the same as young shoot. The

variation of the growth stimulator in MS solid media enriched with 2,4-D: 0; 0.5; and 1.0 mg/l and BAP; 0; 0.5; 1.0, and 1.50 mg/l results in the variation of the culture (Figure 3b). Low intensity of callus is observed when using growth stimulator 2,4-D 0 mg/L and BAP 0.5 mg/L (D_0B_1), where when the culture is enriched with 2,4-D 1.0 mg/L without BAP (D_2B_0) produce high intensity of callus. Callus induction in D_2B_0 is found very good, however it does not produce shoot. The MS medium consisted of 2,4-D 0.5 mg/L without BAP (D_1B_0) could initiate the callus in medium intensity and could directly produce shoot. Therefore, the medium culture that is enriched with 2,4-D 0 mg/L and BAP 0.5 mg/L (D_0B_1) is found effective to initiate the shoot in the shoot tip culture of *Citrus Brastepu*.



Figure 2. The plant appearance of *Citrus nobilis* Var. Brastepu and it's fruit

The growth of Callus in Medium Culture

The culture development in various condition of culture is observed based on the growth of callus, the development of embryosomatic, the intensity of shoot, and the characteristic of the explant in the culture is presented in Table 1. The morphogenesis of the callus is observed from its preparative of the culture (the picture not showed). The microscopic preparation of the callus for meristematic propagation of a shoot is showed in where the shape of the callus is a meristematic cell. The differentiation of a meristem become a nodule of shoot is clearly shown, as could be seen clearly. The callus is dominated by the parenchyma cells that is the active cell in the multiplication process. There is a callus stain in the surface and the nodule that is potentially become regenerated or organogenesis become a plant.

The Development of Embryosomatic of *Citrus nobilis* Var. Brastepu

The formation of embryosomatic of the *Citrus nobilis* Var. Brastepu were conducted by transferring the embryogenic callus into MS medium enriched with 2,4-D and BAP. The variation of the 2,4-D and BAP in the MS medium is sufficient to grow the embryosomatic. The growth and the development of embryosomatic influenced by the variation of growth stimulator in medium is summarized in Table 1. It is known that MS0 medium could not initiate the embryosomatic. The addition of 2,4-D is not also regenerate the embryosomatic until the concentration of the growth stimulator of 1,0 mg/L 2,4-D and 1,0 mg/L BAP, where the best condition is found in D_1B_2 with the texture of embryosomatic is green color, rough texture with nodules.

The Development of Plantlet of *Citrus nobilis* Var. Brastepu

The development of plantlet in a medium culture was categorized as good in all experiments. The explants are consistently grown linearly with the incubation time. The observation for the development of the callus after six weeks incubation time is presented in Figure 3. Extending the incubation time up to seven weeks did not improved the growth intensity of the plantlet. The same phenomena were also found after extending the incubation time up to ten weeks. After the incubation time has been conducted for 12 weeks, the plantlet is develop become a real plants in medium culture, where the root, leaves, the stem and bud are

developed well. The example of for the growth and development of plantlets in medium culture is presented in Figure 4, where the leaves, stem and bud are clearly differentiated become citrus plants. The development of leaves, stem and the roots is clearly showed when the plantlet is taken from medium (Figure 4b) and the variation of the plantlets for a single plant is also presented. With this condition, the plant could be transferred into other media for acclimatization.

Table 1. The development and the characteristic of shoot tip culture of *Citrus nobilis* Var. Brastepu in MS media enriched with 2,4-D and BAP. The number represent the Duncan test (P 0.05).

Experiment Treatment	Callus (g)	Embryosomatics	Shoot	The description and the characteristic of shoot culture
D ₀ B ₀	0.04 a	0.50 a	0.20	Explant with low intensity callus
D ₀ B ₁	0.35 d	1.20 b	0.30	Green explant and big texture
D ₀ B ₂	0.25 c	3.10 d	0.60	Green explant and big texture
D ₀ B ₃	0.16 b	2.30 c	0.50	Green explant and big texture
D ₁ B ₀	1.14 i	9.00 i	1.50	White callus and rough texture
D ₁ B ₁	1.83 l	11.70 j	1.80	Green callus, rough texture
D ₁ B ₂	1.51 k	20.60 l	2.70	Green callus, rough with nodul
D ₁ B ₃	1.37 j	15.50 k	2.10	Green callus, rough with nodul
D ₂ B ₀	0.51 e	4.20 e	0.80	White callus, rough texture
D ₂ B ₁	0.94 h	4.90 f	0.90	Brown callus, rough texture
D ₂ B ₂	0.86 g	7.90 h	1.30	Brown callus, rough texture
D ₂ B ₃	0.62 f	6.10 g	1.10	Green callus, rough texture

D₀ = 2,4-D 0.0 mg/l; D₁ = 2,4-D 0.5 mg/l; D₂ = 2,4-D 1.0 mg/l; B₀ = BAP 0.0 mg/l; B₁ = BAP 0.5 mg/l; B₂ = BAP 1.0 mg/l; B₃ = BAP 3.0 mg/l.



Figure 3. The growth and development of plantlets of local citrus in culture medium enriched with growth stimulator: (a) The plantlet emerged from calli (b) Plenty of plantlet in various sizes.

Isolasi of DNA of *Citrus nobilis* Var. Brastepu and Analysis for CVPD

Isolation of the DNA of from the culture of the *Citrus nobilis* Var. Brastepu have been conducted. An amount of 5 μ L DNA has been taken followed the analysis on gel electrophoresis. The results showed that pure DNA has been produced from the culture which is then be used in PCR analysis for the CVPD screening. The profile of the electrophoresis gel have confimed that all samples are free from CVPD. It is known that from the marker (forward and reverse primers) is specific for *C. Liberibacter asiaticus* which show a single band for the sDNA. It also can be seen from Figure 4. that there are no band for the 24 samples cultures , mean all of them are free from CVPD and the samples with a single band are the CVPD positive.




Figure 4. The electrophoresis of the 24 PCR DNA samples from shoot tip cultures of *Citrus nobilis* Var. Brastepu using primer HLB 65/66

Conclusions

Meristematic shoot tip culture to obtain good quality seedling of citrus (*Citrus nobilis* Var. Brastepu) free from CVPD) have successfully conducted. The best condition by using 0.5 mg/L 2,4-D and 0.5 mg/L BAP (D1B1) in medium culture has been obtained that could initiate the callus, embryosomatic, plantlets, and the plants.

References

- Almeida, R.P.P., Nascimento, F.E., Chau, J., Prado, S.S., Tsai, CW, Lopes, S.A., and Lopes, J.R.S., (2008), Genetic Structure and Biology of *Xylella fastidiosa* Strains Causing Disease in Citrus and Coffee in Brazil, *Applied And Environmental Microbiology*, 74(12): 3690–3701.
- Bove, J.M., (2006)., Huanglongbing: A destructive, newly-emerging, century old disease of citrus. *Journal of Plant Pathology* 88(1): 7-37.
- Carimi, F.; Pasquale, F.D., and Crescimanno, F.G., (1995), Somatic embryogenesis in *Citrus* from styles culture, *Plant Science* 105: 81-86
- Casanova, E.; Trillas, M.I.; Moysset, M. and Vainstein, A., (2005), Influence of *rol* genes in floriculture, *Biotechnology Advances* 23(1): 3-39.
- Chaturvedi, H.C., and Mitra, G.C., (1974), Clonal propagation of citrus from somatic callus culture, *HortScience* 9: 118-120.
- Costa, M.G.C.; Alves, V.S.; Lani, E.R.G; Mosquim, P.R.; Carvalho, C.R., and. Otoni, W. C., (2004), Morphogenic gradients of adventitious bud and shoot regeneration in epicotyl explants of *Citrus*, *Scientia Horticulturae* 100(1-4): 63-74
- Das, A.; Paul, A.K. and Chaudhuri, S., (2000), Micropropagation of sweet orange, *Citrus sinensis* Osbeck. for the development of nucellar seedlings, *Indian Journal of Experimental Biology* 38, 269-272.
- Do Carmo, T.D; Luc, D.J.; Eveillard, S.; Cristina, M.E.; de Jesus, J.WC.; and Takao, Y.P.; Aparecido, L.S.; Beozzo, B.R.; Juliano, A.A.; Saillard, C.; Bove, JM., (2005), Citrus huanglongbing in Sao Paulo State, Brazil: PCR detection of the 'Candidatus' Liberibacter species associated with the disease, *Mol Cell Probes*. 19(3): 173-179.
- Duran-Vila, N.; Ortega, V., and Navarro, L., Morphogenesis and tissue cultures of three citrus species, *Plant Cell and Organ Culture* 16, (1989), 123-133.
- Edriss, M.H., and Burger, D.W., (1984), In Vitro propagation of troyer citrange from epicotyl segment, *Scientia Horticulturae* 23: 159-162.
- Grosser, J.W., and Chandler, J.L., (2000), Somatic hybridization of high yield, cold-hardy and disease resistant parents for citrus rootstock improvement, *Journal Horticultural Science Biotechnology* 75: 641-644.
- Hao, Y.J.; Cheng, Y.J, and Deng, X.X., (2005), Stable maintenance and expression of a foreign gene in transgenic pear shoots retrieved from *in vitro* conservation, *Journal of Plant Physiology* 162(2): 237-243.
- Hidaka, T., and Kajiura, I., (1988), Plantlet differentiation from callus protoplasts induced from *Citrus* embryo, *Scientia Horticulturae* 34(1-2" 85-92
- Hunt, P.W.; Watts, R.A.; (2001), Trevaskis, B.; Llewelyn, D.J.; Burnell, J.; Dennis, E.S. and Peacock, W.J., Expression and evolution of functionally distinct haemoglobin genes in plants, *Plant Molecular Biology* 47: 677-692.

- Kim, J.S., and Wang, N., (2009) Characterization of copy numbers of 16 S DNA and 16 S RNA of *Candidatus Liberibacter asiaticus* and the implication in detection in planta using quantitative PCR. *BMC Research Notes* 2: 37-40.
- Li, W., Song, Q., Bransky, R.H., and Hartung, J.H., (2007), Genetic diversity of citrus bacterial canker pathogens preserved in herbarium specimens, *Proc Natl Acad Sci U S A* 104(47): 18427-18432.
- Maggon, R., and Singh, B.D., Promotion of adventitious bud regeneration by ABA in combination with BAP in epicotyl and hypocotyl explants sweet orange (*Citrus sinensis* L.Osbeck), *Scientia Horticulturae* 63, (1995), 123-128.
- Murashige, T., and Skoog, F., A revised media for rapid growth and bioassay with tobacco tissue culture, *Physiol. Plant.* 15: 473-496.
- Niedz, R.P., and Evens, T.J., (1962), The effects of nitrogen and potassium nutrition on the growth of nonembryogenic and embryogenic tissue of sweet orange (*Citrus sinensis* (L.) Osbeck), *BMC Plant Biol.* 8, (2008), 126-131.
- Nurwahyuni, I., (2003), Kultur jaringan daun jeruk manis (*Citrus nobilis* Brasitepu) untuk mikropropagasi, *Jurnal Sain Indonesia* 24(1): 17-20
- Nurwahyuni, I., (2003), Uji ketahanan kultur jeruk manis (*Citrus nobilis* Brasitepu) terhadap salinitas menuju bibit unggul, *Jurnal Scientia* 3(2): 75-84.
- Nurwahyuni, I., and Elimasni, (2010), *Optimalisasi Teknik In Vitro Untuk Perbanyak Jeruk Berkualitas Baik Sebagai Strategi Biokonservasi Jeruk Lokal Sumatera Utara*, Laporan Penelitian, FMIPA USU Medan.
- Nurwahyuni, I., and Tjondronegoro, P., (1994), Induksi kalus dan regenerasi tanaman *Dioscorea composita* Hemsls, *Hayati* 1: 15-17.
- Prastowo, N.R.; Roshetso, J.M.; Manurung, G.E.S.; Nugraha, E.; Julian, J.M. and Harum, F., (2006), *Teknik pembibitan dan perbanyak vegetatif tanaman buah*. World Agroforestry centre (ICRAF). Bogor.
- Rieger, M., (2006), *Introduction to Fruit Crops*. Hawthorn Food Products Press, NY.
- Sanford M.T. 1992. *Pollination of citrus by honey bees*. UF/IFAS document RFAA092. Available online.
- Su, H. J. (2008), *Production and cultivation of virus free citrus saplings for citrus rehabilitation in Taiwan.*, APCoAB. New Delhi. India.
- Sumaatmaja, G., (2001), Teknik perbanyak tanaman jeruk manis di Kabupaten Bengkulu Selatan. Laporan Teknis PUSLIT-Biologi-LIPI. Jakarta.
- Terol, J., Conesa, A., Colmenero, J.M., Cercos, M., Tadeo, F., Agustí, J., Alós, E., Andres, F., Soler, G., Brumos, J., Iglesias, D.J., Götz, S., Legaz, F., Argout, X., Courtois, B., Ollitrault, P., Dossat, C., Wincker, P., Morillon, R., and Talon, M., (2007), Analysis of 13000 unique *Citrus* clusters associated with fruit quality, production and salinity tolerance, *BMC Genomics*. 8: 31.
- Terol, J., Naranjo, M.A., Ollitrault, P., and Talon, M., (2008), Development of genomic resources for *Citrus clementina*: Characterization of three deep-coverage BAC libraries and analysis of 46,000 BAC end sequences, *BMC Genomics*. 9: 423-429.
- Vu, J.C.V.; Niedz, R.P., and Yelenosky, G., (1995), Activities of sucrose metabolism enzymes in glycerol grown suspension cultures of sweet orange (*Citrus sinensis* L.Osbeck), *Environmental Experimental Botany* 35: 455-463.
- Williamson, J.G., and Jackson, L.K., (1994), *Citrus propagation*. University of Florida, Fact Sheet HS-86, June 1994. <http://ultimatecitrus.com>