

# Pesticide Exposure, Behavior of Farmer, and Activity of Cholinesterase Enzyme in Blood of Fertile Women Farmers

## Pajanan Pestisida, Perilaku Petani, dan Aktivitas Enzim dalam Darah Petani Perempuan Usia Subur

Deni Abdul Rahman\*, Zakianis\*\*, Laila Fitria\*\*

\*Universitas Darussalam Gontor, Ponorogo, Indonesia \*\*Environmental Health Department, Faculty of Public Health, Universitas Indonesia, Depok, Indonesia

DOI: <http://dx.doi.org/10.21109/kesmas.v10i2.879>

### Abstract

Fertile women farmers are risky of suffering decrease of cholinesterase activity due to pesticide exposure. This study aimed to analyze relation between pesticide exposure and the exposure agent to cholinesterase activity of fertile women workers at Kedunguter Village. This study used cross-sectional design on 94 fertile women farmers in 2015. Data was collected by observation, interview and cholinesterase test. Data analysis used chi-square test and analysis results showed a significant relation between pesticide types, working time, the use of gloves, hand-washing behavior to cholinesterase activity of fertile women farmers. Analysis results of this study showed that variable working time had the highest odds ratio (OR) score (OR = 14.072), so the variable working time is the most dominant variable in influencing cholinesterase enzyme. This study suggests that fertile women farmers should work not more than six hours per day.

**Keywords:** Behavior, cholinesterase enzyme, pesticide exposure

### Abstrak

Petani perempuan usia subur berisiko mengalami penurunan aktivitas kolinesterase akibat pajanan pestisida. Penelitian ini bertujuan untuk menganalisis hubungan antara pajanan pestisida dan perilaku pemajan terhadap aktivitas kolinesterase petani perempuan usia subur di Desa Kedunguter. Penelitian ini menggunakan desain potong lintang pada 94 petani perempuan usia subur tahun 2015. Pengumpulan data dilakukan secara observasi, wawancara, dan uji kolinesterase. Analisis data menggunakan uji kai kuadrat dan hasil analisis menunjukkan hubungan signifikan antara jenis pestisida, waktu kerja, penggunaan sarung tangan, perilaku mencuci tangan terhadap aktivitas kolinesterase petani perempuan usia subur. Hasil analisis pada penelitian ini menunjukkan bahwa variabel waktu kerja memiliki nilai *odds ratio* (OR) tertinggi, yaitu OR = 14,072 sehingga waktu kerja merupakan variabel paling dominan dalam memengaruhi enzim ko-

linesterase. Penelitian ini menyarankan agar petani perempuan usia subur tidak bekerja lebih dari enam jam per hari.

**Kata kunci:** Perilaku, enzim kolinesterase, pajanan pestisida

### Introduction

Brebes District is the largest red onion-producing region in Central Java.<sup>1</sup> The higher production of red onion, the more pesticides used in Brebes District. Pesticides most commonly used in Brebes District are organophosphate and carbamate insecticides.<sup>2</sup> Organophosphates and carbamates are substance that may inhibit or inactivate cholinesterase enzyme activity.<sup>3</sup>

Health agency has conducted an observation of cholinesterase enzyme activity in blood of onion farmers in Brebes District every year. Results of the observation within the last five years showed that proportion of the highest incidence of cholinesterase disorder occurred in 2012 was 56%. The observation in 2012 was conducted in 10 villages over Brebes District and the observation results showed that Kedunguter Village in Brebes Subdistrict was the village with the highest incidence of cholinesterase enzyme disorder worth 82%. Therefore, Kedunguter Village was selected as the location for this study.<sup>4</sup>

According to the prior study, any disorder/decrease of cholinesterase enzyme activity was found among fertile

Correspondence: Deni Abdul Rahman, Universitas Darussalam Gontor, Siman KM 6 Street Ponorogo District, East Java, Indonesia, Phone: - , e-mail: [deni.abrah.7@gmail.com](mailto:deni.abrah.7@gmail.com)/[deni@unida.gontor.ac.id](mailto:deni@unida.gontor.ac.id)

women around farming area.<sup>5</sup> Fertile women are a vulnerable group due to pesticide exposure, considering effect of pesticides is substance that may influence establishment and growth of embryo.<sup>3,6</sup> This statement is supported by the prior study conducted in Poland. Based on that study, there was a significant relation between pesticide exposure among mothers and incidence of low birth-weight (LBW) and premature pregnancy. Disorder on pregnancy process may cause harm effects for child growth.<sup>7</sup> Based on Government Regulation (*Peraturan Pemerintah*) 66 Article 44 of The Year 2014, one of actions to protect people from pesticide dangers is analyzing harmful risks of pesticide exposure impacts.<sup>8</sup> Risk analysis can be conducted by measuring cholinesterase enzyme activity of fertile women farmers.<sup>2</sup>

This study is not the first study conducted among fertile women at work scope of Brebes District Health Agency. The similar kind of study was already conducted by several researchers who observed relation between fertile women living in farming area and cholinesterase enzyme activity. Differ from previous studies, all respondents in this study were fertile women working as farmers, so all the respondents in this study were a risky group and observation was conducted in obtaining information concerning behavior of fertile women farmers. This study aimed to analyze relation between pesticide exposure and behavior of the exposure agent to cholinesterase enzyme activity of fertile women farmers and to determine dominant factors influencing the cholinesterase enzyme activity of fertile women farmers at Kedunguter Village, Brebes Subdistrict, Brebes District in 2015.

**Method**

This study used cross-sectional design. Cross-sectional design is study in aim to observe relation between risk factors and effects at the same time. Location of study was at Kedunguter Village, Brebes Subdistrict, Brebes District. This study was conducted on February 20<sup>th</sup> – May 20<sup>th</sup> 2015. Population was all fertile women working as active farmers because they always did red onion farming activity at field, but did not do pesticide spraying activity and they lived at Kedunguter Village, Brebes Subdistrict, Brebes District.

Sample of this study was as many as to 94 respondents obtained by using proportion estimating formula. Samples were fertile women at Kedunguter Village, Brebes Subdistrict, Brebes District selected by using random sampling method. Data was received from direct observation in the location of study namely interviewing respondents as conducted within one week before cholinesterase enzyme examined. The examination of cholinesterase enzyme activity that used Tintometer Kit Lovibond device was conducted by Brebes District Health Agency with doctors around the nearest primary

health care while this study was being conducted, or with laboratory officer appointed by Brebes District Health Agency.<sup>4,9</sup> Results of any relation found in this study was assessed by considering results of p value, odds ratio (OR) and confidence interval (CI).

**Results**

**Cholinesterase Enzyme Activity**

Decrease of cholinesterase enzyme activity was the impact emerged by excessive pesticide exposure. Being fertile woman farmer is such a work vulnerable to pesticide exposure. Table 1 presented data of results of cholinesterase test conducted among 94 fertile women farmers at Kedunguter Village in 2015.

In this study, 100% of fertile women farmers at Kedunguter Village had normal cholinesterase enzyme activity because of not having cholinesterase enzyme activity  $\leq 75\%$  based on cut off point. Therefore, it needed determination of new cut off point in aim to ease statistical analysis. Before determining the new cut off point, data normality test (*kolmogorov smirnov*) was early conducted to ensure normality of data distribution.

Results of data normality test on cholinesterase enzyme activity showed p value  $< \alpha$  (0.05), so the distribution form was abnormal. Based on this weakness, a new cut off point was determined based on statistical approach by using quartile value. The middle value of the first quartile would be used as the new cut off point for cholinesterase activity in blood of fertile women farmers because the first quartile value was the value closest to the threshold of pesticide poisoning level. Table 2 presented category of cholinesterase activity based on result of cut off point. Table 2 showed most fertile women farmers who had activity  $> 87.5$ .

**Table 1. Results of Cholinesterase Enzyme Activity Test and Data Normality Test on Cholinesterase Enzyme Activity of Fertile Women Farmers**

Cholinesterase Enzyme Activity	n	%	Kolmogorov-Smirnov			Q1	Q2	Q3
			Statistic	Df	p value			
75%	6	6.4	0.372	94	0.000	87.5	100	100
87.5%	32	34.0						
100%	56	59.6						
<b>Total</b>	<b>94</b>	<b>100.0</b>						

**Table 2. Distribution of Frequency of Cholinesterase Enzyme Activity in Fertile Women Farmers**

Category of Cholinesterase Enzyme Activity of Fertile Women Farmers	n	%
$> 87.5$	38	40.4
$\leq 87.5$	56	59.6
<b>Total</b>	<b>94</b>	<b>100</b>

**Pesticide Exposure**

Pesticide exposure got by fertile women farmers was the result from pesticide spraying activity done by men farmers. Fertile women farmers were active farmers who always did farming every day, but did not do pesticide spraying. The pesticide spraying activity was only performed by men farmers, so information regarding pesticide types was obtained from the men farmers. The information regarding pesticide types was obtained from observation conducted among 82 men farmers (pesticide spraying agent). Men farmers always used many types of pesticides and mixed two to three types of pesticides in spraying. Table 3 presented results of observation concerning pesticide types used by men farmers.

The information regarding types of active materials

**Table 3. Frequency of Types of Active Materials Used as Pesticide Mixture by 82 Farmers Spraying Pesticides**

Type	Active Materials	n	%
Organophosphate	Profenofos	31	42
	Phenthoate	23	32
	Chlorpyrifos	19	26
<b>Total</b>		<b>73</b>	<b>100</b>
Non-organophosphate	Carbosulfan	23	17
	Cypermethrin	30	22
	Carbosulfan	23	17
	Chlorfenapyr	33	25
	Lufenuron	25	19
<b>Total</b>		<b>134</b>	<b>100</b>

on Table 3 showed category of pesticide types received by fertile women farmers from pesticide types used by the spraying agent. Type of pesticide exposure among fertile women farmers was determined by considering the nearest location of the spraying agent with fertile women farmers while the observation conducted.

Table 4 presented relations of independent variables (pesticide exposure and the behavior of the exposing agent) and dependent variable (cholinesterase activity). Data presented on Table 4 showed a significant relation between pesticide types, working time, the use of gloves and hand-washing behavior to cholinesterase enzyme activity of fertile women farmers according to OR score.

Final model of this study showed that working time was the dominant factor influencing cholinesterase enzyme activity with OR score = 14.072 (Table 5).

**Discussion**

Results of cholinesterase enzyme activity test on fertile women farmers at Kedunguter Village showed 100% normal because the spraying farmers used more pesticide materials not from organophosphate category. Organophosphate and carbamate entering to the body tend to be bounded with cholinesterase enzyme that caused inactivation of cholinesterase enzyme. Such matter caused acetylcholine piled on nerve and occurrence of overstimulus on receptor nerve.<sup>10</sup> Most spraying farmers used organophosphate pesticide only for additional mixing material, so the dose used was little. Although

**Table 4. Analysis of Relation between Pesticide Exposure and Cholinesterase Activity in Blood of Fertile Women Farmers**

Variable	Category	Cholinesterase Activity						OR (95%CI)	p value
		≤ 87.5%		> 87.5 %		Total			
		n	%	N	%	N	%		
Pesticide exposure									
Pesticide types	Organophosphate	28	47.5	31	52.5	59	100	2.258	0.113
	Non-organophosphate	10	28.6	25	71.4	35	100	(0.924 - 5.520)	
Working period	> 13.5 years	19	40.4	28	59.6	47	100	1.000	1.000
	≤ 13.5 years	19	40.4	28	59.6	47	100	(0.439 - 2.279)	
Working time	> 6 hours/day	21	80.8	5	19.2	26	100	12.6	0.000
	≤ 6 hours/day	17	25.0	51	75.4	68	100	(4.115 - 38.585)	
Behavior of exposure act									
Use of mask	Not using mask	37	41.6	52	58.4	89	100	2.846	0.645
	Using mask	1	20	4	80	5	100	(0.306 - 26.508)	
Wearing long sleeve shirt	Not wearing long sleeve shirt	2	100	0	0	2	100	0	0.161
	Wearing long sleeve shirt	36	39.1	56	60.9	92	100	(0-0)	
Use of gloves	Not using gloves	13	56.05	10	43.5	23	100	2.392	0.117
	Using gloves	25	35.2	46	64.8	71	100	(0.918 - 6.232)	
Hand-washing behavior	Not washing hand before eating	34	46.6	39	53.4	73	100	3.705	0.026
	Washing hand before eating	4	19	17	82	21	100	(1.136 - 12.086)	
Bathing behavior	Not immediately bathing and dressing up after going home from rice field	2	66.7	1	33.3	3	100	3.056	0.563
	Immediately bathing and dressing up after going home from rice field	36	39.6	55	60.4	91	100	(0.267 - 34.950)	
<b>Total</b>		<b>38</b>	<b>40.4</b>	<b>56</b>	<b>59.6</b>	<b>94</b>	<b>100</b>		

Table 5. Final Model of Multivariate Analysis (Multiple Logistic Regression Analysis)

Variable	B	p value	OR	CI 95%	
				Lower	Upper
Pesticide types	0.961	0.089	2.614	0.863	7.915
Working time	2.644	0.000	14.072	4.113	48.149
Use of gloves	0.776	0.199	2.173	0.665	7.101
Hand washing behavior	1.469	0.053	4.346	0.979	19.305
Constant	-8.553	0.000	0.000		

organophosphate was only used as additional mixing material while spraying pesticides, some fertile women farmers were found suffering the decrease of cholinesterase enzyme activity. There were 38 fertile women farmers who had cholinesterase enzyme activity  $\leq 87.5\%$ .

Organophosphate is the main agent in the decrease of cholinesterase enzyme activity.<sup>10,11</sup> Bivariate analysis showed significant relation between pesticide type used by the spraying farmers and cholinesterase enzyme according to OR score. Meanwhile, multivariate analysis showed that pesticide type used by farmers would contribute to cholinesterase enzyme activity. Result of this study was aligned with previous study obtaining relation between pesticide types used by farmers and cholinesterase enzyme activity.<sup>12</sup> Information about pesticide types was obtained by conducting observation among men farmers, land owners, or husbands of fertile women farmers who were spraying pesticides. After the information was obtained, such information would be categorized based on active materials and the pesticide types.

The longer a person worked by using pesticides, it would be aligned with effects of the decrease of cholinesterase enzyme activity.<sup>12</sup> However, such case was not found in this study. Based on bivariate analysis results, no significant relation was found between working period and cholinesterase enzyme activity. Result of this study was aligned with previous study which did not find out significant relation between working period and cholinesterase enzyme activity.<sup>13-15</sup> According to bivariate analysis, 50% of fertile women farmers had working period  $\geq 13.5$  years, but this did not give any influence to farmer's cholinesterase enzyme activity due to little dose got by fertile women farmers. Based on their activities, fertile women farmers would get pesticide exposure from irrigation water channel in rice field and pesticide residue on plants. Study conducted by Taufik & Yosmaniar,<sup>16</sup> showed that Brebes District irrigation channel and plantation was polluted by organophosphate pesticide residue worth 2.7 – 3.2 g/L.

The longer a person in contact with pesticides every day, the more decreasing cholinesterase activity would

be. Therefore, the maximum threshold to contact with pesticides should be no more than five hours per day, even Environmental Protection Agency (EPA) has smaller maximum threshold that is four hours per day to contact with pesticides.<sup>17,18</sup> Results of bivariate analysis showed significant relation between farmer's working time per day and cholinesterase enzyme activity, meanwhile multivariate analysis showed that working time was the dominant factor in influencing cholinesterase enzyme activity in blood of fertile women farmers. This was in accordance with the previous study that found out significant relation between working time and cholinesterase enzyme activity in blood.<sup>19</sup>

This study found out fertile women farmers who worked more than six hours every day. They did additional activities, such as watering plants and harvesting. Based on reference, ability of organophosphate to enter and start influencing cholinesterase enzyme may occur in relatively short time.<sup>11</sup> So that, from the analysis results of this study, fertile women farmers are hoped not to do farming activities more than six hours every day. It will definitely decrease productivity of fertile women farmers, so to maintain the productivity, fertile women farmers can do any other additional activities, such as participating at entrepreneurship activities that can maintain and increase their productivity. Therefore, government should provide any entrepreneurship training for fertile women farmers to avoid quite long pesticide exposure effects.

The use of mask takes an important role inhibiting the entering of pesticides to human's body through inhalation.<sup>20</sup> Disobedience to the use of self-protector (*alat pelindung diri/APD*) significantly related to cholinesterase on farmers.<sup>21</sup> However, this study found out different results. Based on bivariate analysis, OR score = 2.846 (0.306 – 26.508). Because OR = 0.306 – 26.508 passed 1, this showed that no significant relation was found between the use of nose and mouth-covering mask and cholinesterase enzyme activity. Pesticides would be easier to be inhaled by fertile women farmers who did not use such mask while working. The pesticides inhaled by farmers can directly enter to blood circulation system in short time namely in second to minute, so it is regrettable if there are still many fertile women farmers who never use mask while working at the field.<sup>11</sup> Most fertile women farmers do not use nose-covering mask. Therefore, fertile women farmers always feel such tightness while using the mask. Brebes District Health Agency should hold any counseling and grant operational assistance to fertile women farmers in providing masks worthy used during farming activities.

Fertile women farmers should wear long sleeve shirt to avoid direct exposure of sprayed pesticides as this action is the early step to prevent the sprayed pesticides entering through the skin.<sup>22</sup> Result like variable self-pro-

tektor use was previously found on variable self-protector use in form of long sleeve shirt. Bivariate analysis in this study showed no significant relation between the use of long sleeve shirt and cholinesterase enzyme activity. Result of this study was aligned with the prior study which did not obtain significant relation between the use of long sleeve shirt and cholinesterase enzyme activity in farmer's blood.<sup>23</sup>

While applying pesticides, farmers should always use gloves as self-protector. The previous study had explained that disobedience to the use of self-protector would increase risk of decrease of cholinesterase activity in blood.<sup>21,24</sup> The result of bivariate analysis showed any significant relation between the use of gloves and cholinesterase enzyme activity of fertile women farmers based on OR score, meanwhile multivariate analysis showed that variable the use of gloves was the factor contributing to cholinesterase enzyme activity. Most fertile women farmers had used gloves while working at rice field, but all the gloves used by the farmers were made of fabric. Gloves made of fabric are easily traversed by liquid into the skin, so farmers should replace it by gloves made of rubber material which is liquid proof.

Washing hands before eating is self-protection in form of personal hygiene behavior to avoid exposure of any substance to enter through mouth.<sup>20,25</sup> Hands unwashed before eating will bring pesticides attached on hands, so this can cause pesticides contaminate food and enter through alimentary canal. Time needed for pesticides to enter into the body through oral may happen fast from 30 to 90 minutes.<sup>11</sup> Therefore, fertile women farmers should always keep their hands clean before eating whether at rice field or not. Bivariate analysis showed any significant relation between hand-washing behavior of farmers and cholinesterase enzyme activity, meanwhile multivariate analysis showed that hand-washing activity was the factor contributing to cholinesterase enzyme activity of fertile women farmers. Such results were aligned with the prior study that found significant relation between hand-washing behavior and cholinesterase enzyme activity in blood.<sup>26</sup>

Similar to the previous variable, variable bathing behavior was one of behaviors that belong to personal hygiene. However, bivariate analysis showed no any significant relation between farmer's bathing behavior and cholinesterase enzyme activity in blood of fertile women farmers. Based on this study, most fertile women farmers had good behavior by taking a bath after going home from farming. This was an act of fertile women farmers to remove any pesticides attached on the body. This was aligned with the prior results of study that did not found significant relation between personal hygiene and cholinesterase enzyme activity in blood.<sup>15</sup> Bathing is one of acts to remove pesticide exposure attached on the

body, therefore good practice of bathing can be taken after using pesticides, such as spraying, mixing or doing other farming activities that contain activities using pesticides.<sup>18</sup>

### Conclusion

Based on analysis results, there are four variables related to the decrease of cholinesterase enzyme activity of fertile women farmers at Kedunguter Village namely pesticide types, working time, glove using and hand-washing behaviors. Variable working time is the most dominant variable to the decrease of cholinesterase activity  $\leq 87.5\%$ .

### Recommendation

Working time is the dominant variable influencing cholinesterase enzyme activity in blood of fertile women farmers, so health agency and agriculture agency should provide counseling for the farmers to not spend time of farming more than six hours per day. The working time for six hours per day among fertile women farmers can decrease their work productivity and income. An act to maintain and increase productivity of fertile women farmers can be performed by any multisectoral activities between health agency, agriculture agency and other social institutions in Brebes District. Activities that can be held are entrepreneurship training or opening job opportunities that can maintain productivity as well as increase income of fertile women farmers at Kedunguter Village.

### References

1. Sayaka B, Supriyatna Y. Kemitraan pemasaran bawang merah di Kabupaten Brebes, Jawa Tengah. Seminar nasional peningkatan daya saing agribisnis berorientasi kesejahteraan petani. Bogor: Pusat Analisis Sosial Ekonomi dan Kebijakan Pertanian, Badan Penelitian dan Pengembangan Pertanian Departemen Pertanian; 2009 [cited 2014 Jun 5]. Available from: [http://pse.litbang.pertanian.go.id/ind/pdffiles/MKP\\_C2.pdf](http://pse.litbang.pertanian.go.id/ind/pdffiles/MKP_C2.pdf).
2. Marinajati D, Endah N, Suhartono. Hubungan riwayat paparan pestisida dengan profil darah pada wanita usia subur di daerah pertanian cabai dan bawang merah. *Jurnal Kesehatan Lingkungan Indonesia*. 2012; 11 (1): 61-7.
3. Society of Toxicology. *The toxicologist*. San Antonio (Texas): Oxford University; 2013.
4. Departemen Kesehatan Lingkungan Dinas Kesehatan Brebes. Rekapitulasi hasil kegiatan pemeriksaan cholinestrerase darah petani yang kontak dengan pestisida tahun 2012. Brebes: Dinas Kesehatan Brebes; 2013.
5. Siwiendrayanti A, Suhartono, Endah N. Hubungan riwayat pajanan pestisida dengan kejadian gangguan fungsi hati (studi pada wanita usia subur di Kecamatan Kersana Kabupaten Brebes). *Jurnal Kesehatan Lingkungan Indonesia*. 2012; 11 (1): 9-14.
6. Damanhuri E. *Pegolahan bahan berbahaya dan beracun (B3)*. Diktat Pengelolaan B3. 2009-2010. Bandung: Institut Teknologi Bandung;

- 2010.
7. Dabrowski S, Hanke W, Polanska K, Makowiec-Dabrowska T, Sobala W. Pesticide exposure and birthweight: an epidemiological study in Central Poland. *International Journal of Occupational Medicine and Environmental Health* [serial on internet]. 2003 [cited 2015 Jan 5]; 16(1): 31–39. Available From: <http://test.imp.lodz.pl/upload/oficy-na/artkuly/pdf/full/Dab3-01-05.pdf>.
  8. Presiden Republik Indonesia. Peraturan Pemerintah Republik Indonesia. No 66. tahun 2014 tentang kesehatan lingkungan. Jakarta: Kementerian Sekretariat Negara Republik Indonesia; 2014.
  9. The tintometer ltd. AF267 cholinesterase test kit [monograph on the Internet]. 2015 [cited 2015 Jan 5]. Available from: <http://www.lovi-bondcolour.com/instrument/af267-cholinesterase-test-kit>.
  10. World Health Organization. Clinical management of acute pesticide intoxication: prevention of suicidal behaviours. Geneva, Switzerland: World Health Organization; 2010.
  11. Agency for Toxic Substances and Disease Registry. Agency for toxic substances and disease registry case studies in environmental medicine cholinesterase inhibitors: including pesticides and chemical warfare nerve agents. Agency for Toxic Substances and Disease Registry. 2010 [cited 2014 Desember 28]. Available from: <http://www.atsdr.cdc.gov/csem/cholinesterase/docs/cholinesterase.pdf>
  12. Marsaulina I, Wahyuni AS. Faktor- faktor yang berhubungan dengan keracunan pestisida pada petani hortikultura di Kecamatan Jorlang Hataran Kabupaten Simalungun Tahun 2005. *Media Litbang Kesehatan*. 2007; 17 (1): 18-25.
  13. Rustia HN, Wispriyono B, Susanna D, Luthfiah FN. Lama pajanan organophospat terhadap penurunan aktivitas enzim kolineterase dalam darah petani sayur. *Makara Seri Kesehatan*. 2010; 14 (2): 95-101.
  14. Mahmudah M, Wahyuningsih NE, Setyani O. Kejadian keracunan pestisida pada istri petani bawang merah di Desa Kedunguter Kecamatan Brebes Kabupaten Brebes. *Media Kesehatan Masyarakat Indonesia*. 2012; 11 (1): 65-70.
  15. Budiawan AR. Faktor risiko yang berhubungan dengan cholinesterase pada petani bawang merah di Ngurensiti Pati. *Jurnal Kemas UNNES*. 2014; 3 (1): 1-11.
  16. Taufik I, Yosmaniar. Pencemaran pestisida pada lahan perikanan di daerah Karawang - Jawa Barat. *Prosiding Seminar Nasional Limnologi* V. 2010 [cited 2015 Mei 5]. Available from: [http://limnologi.lipi.go.id/katalog/index.php/searchkatalog/downloadDataById/360/52\\_Pencemaran\\_pestisida\\_pada\\_lahan\\_perikanan\\_Imam\\_Taufik.pdf](http://limnologi.lipi.go.id/katalog/index.php/searchkatalog/downloadDataById/360/52_Pencemaran_pestisida_pada_lahan_perikanan_Imam_Taufik.pdf)
  17. Kementerian Pertanian Republik Indonesia. Peraturan Menteri Pertanian nomor 76/permentan/OT.140/12/2012. Syarat dan tata cara penetapan produk unggulan hortikultura. Jakarta: Kementerian Pertanian Republik Indonesia; 2012.
  18. US Environmental Protection Agency. Recognition and management of pesticide poisonings. Washington DC: Environmental Protection Agency; 2015.
  19. Prado-Lu JL. Pesticide exposure, risk factors and health problems among cutflower farmers: a cross sectional study. Philippines: *Journal of Occupational Medicine and Toxicology*. 2007; 2 (9): 1-8.
  20. US Environmental Protection Agency. Citizen's guide to pest control and pesticide safety. United States: US Environmental Protection Agency; 2005.
  21. Ming Ye, Beach J, Martin JW, Senthilselvan A. Occupational pesticide exposures and respiratory health. *International Journal of Environmental Research and Public Health*. 2013; 10: 6442-71.
  22. OHS The University Of Queensland. Health surveillance guideline for users of organophosphate pesticides [monograph on the Internet]. University Of Queensland. 2012 [cited 2015 Jan 4]. Available from: <http://www.uq.edu.au/ohs/HEALTH/ophealthsurveillance.pdf>
  23. Kartika Y. Faktor risiko yang berkaitan dengan kejadian keracunan pestisida pada petani penyemprot tanaman bawang merah di Desa Sengon Kecamatan Tanjung Kabupaten Brebes [undergraduate thesis]. Semarang: Universitas Negeri Semarang; 2012.
  24. WorkSafeBc. Working safely with OPs (Organophosphate Insecticides) [monograph on the Internet]. WorkSafeBc Canada. 2012 [cited 2015 Feb 5]. Available from: [http://www.worksafebc.com/publications/health\\_and\\_safety/by\\_topic/assets/pdf/organophosphates.pdf](http://www.worksafebc.com/publications/health_and_safety/by_topic/assets/pdf/organophosphates.pdf).
  25. Djojosumarto P. Teknik aplikasi pestisida pertanian. Yogyakarta: Kanisius Media; 2008.
  26. Suwastika IN. Faktor lingkungan, higiene perorangan, perilaku penyemprot dan tingkat keracunan pestisida petani jeruk di Kabupaten Timor Tengah Selatan [postgraduate thesis]. Yogyakarta: Universitas Gadjah Mada; 2009.