

## **Evaluation on Symptoms and Self-Treatment due to Radiotherapy and Chemotherapy Based on Different Cancer Stage**

**Yesiana Dwi Wahyu Werdani<sup>1\*</sup>, Arief Widya Prasetya<sup>2</sup>**

<sup>1</sup> *Widya Mandala Catholic University Surabaya, East Java, Indonesia*

<sup>2</sup> *STIKES Katolik St. Vincentius a Paulo Surabaya.*

*ywerdani@yahoo.com*

### **ABSTRACT**

Cancer modality therapy can minimize metastatic, but it causes decrease in body functions and interfere the daily needs fulfilled. This study was to identify symptoms and self-treatment due to radiotherapy and chemotherapy based on the different cancer stage. This was mixed method design with sequential explanatory approach. Samples were 30 cancer patients taken by purposive sampling at Indonesian Cancer Foundation East Java Branch Surabaya. The instrument was a questionnaire. Fatigue was the most symptoms complained by all patients. Pain, hair loss, constipation were dominated in cancer stage 4, nausea-vomiting, insomnia were dominated in cancer stage 3. The majority self-treatment of fatigue was yoga exercise. Pain and nausea-vomiting were treated by medication and relaxation. Insomnia was treated by mild massage. Constipation was treated by consuming the high fiber foods. Radiotherapy and chemotherapy affect the disturbance of metabolism that cause physical symptoms disorders. Self-treatment by the patient could reduce the symptoms effectively.

**Keywords:** Radiotherapy, Chemotherapy, Cancer Stage, Self-Treatment

Received December, 25, 2019; Revised January 24, 2020; Accepted February 15, 2020



STRADA Jurnal Ilmiah Kesehatan, its website, and the articles published there in are licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.

## INTRODUCTION

Cancer is tumor cells which has the ability to replicate rapidly and invade the normal body tissues (Bunz, 2008). Cancer can metastatic to all of organs depending on the stage of cancer. Prevention of metastatic can be done through radiotherapy and chemotherapy. It is expected to has a longer survival. In prior studies of women with breast cancer post-chemotherapy found that the survival rates of patients increased reaching 5 years (92.1%) and 10 years (81.9%) (Rossi et al., 2015). The effectiveness of radiotherapy and chemotherapy in prolonging survival was followed by various effects on physical conditions. It will lead to disruption of the patient's daily activities and if it lasts longer can worsen the quality of life of the patient. In the prior studies only declared about the various physical effects post-chemotherapy and radiotherapy separately. No research identify these effect and the beneficial treatment by patient to reduce it based on the cancer stage.

DI Yogyakarta is the highest prevalence cancer in Indonesia reached 4.9‰. The most are women (2.9‰) and in urban areas (2.06‰). There are many cancer therapy, the most is surgery reached 61.8%, chemotherapy 24.9%, radiation 17.3% and other therapies 24.1% (Badan Penelitian dan Pengembangan Kesehatan, 2018).

Research on 100 cancer patients found that the majority effects of chemotherapy is a weakness 95% and fatigue 90% (Aslam et al., 2014). Another research reported that 15 patients performed radiotherapy, majority complained were totally exhausted, pain like skin burning and itching (Schnur, Ouellette, Bovbjerg, & Montgomery, 2009). Radiotherapy and chemotherapy trigger the occurrence of oxidative stress not only in the target tissue, but also on healthy tissue. It causes the toxicity and organ dysfunction (Gilliam & St. Clair, 2011). Exposure to oxidative stress on radiotherapy and chemotherapy to the muscles can cause weakness and progressively increase the occurrence of fatigue (Powers & Jackson, 2008). Oxidative stress can lead to apoptosis, metabolic disorders, neuroinflammation, metabolic disturbances and neuronal failure (Areti, Yerra, Naidu, & Kumar, 2014). Target cell damage due to radiation on the tissues can cause DNA damage (Baskar, Dai, Wenlong, Yeo, & Yeoh, 2014).

Therefore the physical symptoms due to chemotherapy and radiotherapy is very important to be recognized by patients earlier and empower patients to reduce symptoms independently is essential to improve the patient's adaptation process. This study was to identify the physical symptoms and self-treatment due to radiotherapy and chemotherapy based on the cancer stage.

## METHODS

This was mixed method design with sequential explanatory approach. Sample was cancer patients with radiotherapy and chemotherapy program at Indonesian Cancer Foundation East Java Branch Surabaya which amounted to 30 people, taken by purposive sampling technique. The questionnaire as an instrument was made by researchers based on theories about the effects of chemotherapy and radiotherapy. It consists of closed-ended questions that describe a wide range of physical complaints in post-chemotherapy and radiotherapy. The instrument has been tested the validity and reliability test, the result of the correlation coefficient was 0.872 and cronbach's alpha result was 0.923. It means that the instrument is valid and reliable. All participants were given an explanation of the purposes, benefits and risks of this study, and who agreed to sign a voluntary and non-compulsory informed consent sheet. After the participants filling out the questionnaire, it continued with deep interviewed about the self-treatment to reduce the symptoms. After the data collected, it

was compare among the symptoms and self-treatment based on cancer stage and calculate the percentage. Interviewed result was presented as a qualitative data.

## RESULTS

Table 1. Demographic Data of Participants

Demographic Data	Categories	Frequency (Person)	Percentage (%)
Age	26 – 35 years old	3	10
	36 – 45 years old	4	13.3
	46 – 55 years old	12	40
	55 – 65 years old	9	30
	> 65 years old	2	6.7
Sex	Female	22	73.3
	Male	8	26.7
Type of cancer	Breast cancer	9	30
	Cervixal cancer	6	20
	Nasopharing cancer	8	26.7
	Colon caner	2	6.7
	Ovary cancer	2	6.7
	Lung cancer	1	3.3
	Lymphoma Hodgkin	1	3.3
	Rectum cancer	1	3.3
Cancer stage	I	4	13.3
	II	6	20
	III	12	40
	IV	8	26.7
Time of diagnosed with cancer	< 1 years	11	36.7
	1 - < 2 years	12	40
	2 - < 3 years	2	6.7
	> 3 years	5	16.6

Table 1 shown the majority participants were female (73.3%) with average of age 46 – 55 years old (40%). The most type of cancer was breast cancer (30%), based on cancer stage was III (40%) and majority time of diagnosed with cancer was 1 - < 2 years (40%).




Figure. 1. The Symptoms due to Radiotherapy and Chemotherapy Based on Cancer Stage

Graph as shown in fig. 1 that found six symptoms due to radiotherapy and chemotherapy, but fatigue was found in all of stage of cancer. In stage 1 no symptom about hair loss and constipation. In stage 2 – 4 all symptoms was found. Pain, hair loss and constipation symptoms were dominated in cancer stage 4, whereas nausea/ vomiting and insomnia were dominated in cancer stage 3.

Table 2. Description of details characteristic symptoms due to radiotherapy and chemotherapy based on cancer stage (section 1)

Stage of Cancer	Fatigue		Pain		Nausea	
	Weakness & lack of energy (%)	Fatigue disturbing of daily activity (%)	Hand, feet & back cramps (%)	Abdominal & back lancinating (%)	When smell of food sometimes followed by vomiting (%)	Any time sometimes followed by vomiting (%)
Stage 1	100	0	75	25	100	0
Stage 2	66,7	33,3	75	25	100	0
Stage 3	25	75	60	40	10	90
Stage 4	0	100	28,6	71,4	14,3	85,7

Table 2 shown that fatigue symptom identified two details characteristics. The first was weak and lack of energy, it was dominated in cancer stage 1 and the second fatigue disturbing of daily activity fulfilling was dominated in cancer stage 4. In Pain symptom participants felt hand, feet and back cramps was dominated in cancer stage 3 and abdominal & back lancinating was dominated in cancer stage 4. Participants felt nausea when smell of food and sometimes followed by vomiting was dominated in cancer stage 1 and 2. Nausea feel anytime sometimes followed by vomiting was dominated in stage 3.

Table 3. Description of details characteristic of symptom due to radiotherapy and chemotherapy based on cancer stage (section 2)

Stage of Cancer	Insomnia		Hair Loss			Constipation		
	Trouble getting started sleeping & Woke up early (%)	Woke up at night and could not sleep anymore (%)	Slight (%)	Moderate (%)	Severe (%)	Less than twice a week (%)	Straining during defecation (%)	Hard lump of feces (%)
Stage 1	100	0	0	0	0	0	0	0
Stage 2	33	66,7	100	0	0	100	100	100
Stage 3	66,7	33,3	0	85,7	14,3	100	100	100
Stage 4	0	100	0	66,6	33,4	100	100	100

Table 3 shown that participants got trouble getting started sleeping & woke up early were dominated by cancer stage 3, and woke up at night and could not sleep anymore were dominated in cancer stage 4. In hair loss symptom the most moderate hair loss was dominated in cancer stage 3 and severe hair loss (alopecia) was dominated in cancer stage 4.

In constipation symptom all of stage of cancer felt straining during defecation and hard lump of faeces except cancer stage 1.

Table 4. Description of details self-treatment to reduce the symptoms (section 1)

Stage of Cancer	Fatigue		Pain		Nausea/ Vomiting	
	Resting (%)	Yoga Exercise (%)	Analgesic drug (%)	Combination of analgesic drug & relaxation (%)	Antiemetic drug (%)	Combination of antiemetic drug & relaxation (%)
Stage 1	25	75	50	50	0	100
Stage 2	50	50	33.3	66.4	0	100
Stage 3	33,3	66,7	30	70	30	70
Stage 4	62,5	37,5	25	75	42,9	57,1

Table 4 shown that in fatigue symptoms, participants relieve it by resting was dominated in cancer stage 4, but yoga exercise was dominated in cancer stage 1. In pain symptom the majority of cancer stage 3 relieve by analgesic drug consume, but treatment by combination of analgesic drug & relaxation (deep breathing and distraction) were dominated by cancer stage 4. Majority participants in cancer stage 4 relieve nausea/ vomiting by antiemetic drug, but in cancer stage 1 and 2 relieve by combination of antiemetic drug & relaxation (deep breathing).

Table 5. Description of details self treatment to reduce the symptoms (section 2)

Stage of Cancer	Insomnia		Hair Loss		Constipation
	Mild massage (%)	Sleeping pills (%)	No Treatment (%)	Using hair tonic (%)	High fiber food consuming (%)
Stage 1	100	0	0	0	0
Stage 2	100	0	100	0	100
Stage 3	66,7	33,3	42,9	57,1	100
Stage 4	40	60	50	50	100

Table 5 shown that mild massage on head, back, hand and feet were dominated in cancer stage 1 and 2 to relieve insomnia, whereas cancer stage 4 relieve insomnia by sleeping pills. In symptom of hair loss the majority of participants in cancer stage 2 do no treatment, and using hair tonic to stimulate the hair growing were used by cancer stage 3. In symptoms of constipation participants in cancer stage 2 – 3 was relieved by high fiber foods consuming such as vegetables, fruits and drink amount of water.

## DISCUSSION

### Symptoms and its characteristics

The most symptoms after chemotherapy and radiotherapy was fatigue, it occurred in all patients with various stages of cancer. The details characteristic identified were weakness, lack of energy and fatigue disturbing of daily activity. The participants have limitations in the

activity and cause the high levels of dependence on others. One of research on 100 cancer patients reported that weakness (95%) and fatigue (90%) after chemotherapy (Aslam et al., 2014). Fatigue (cancer related fatigue) is a subjective complaint against the onset of physical fatigue. Post radiotherapy or chemotherapy majority of respondents got severe fatigue (Kummer, Catuogno, Perseus, Bloch, & Baumann, 2013), (Bock, Hanisch, Matthes, & Zänker, 2014), (Werdani, 2017). Chemotherapy, surgery, and radiotherapy can lead to increased plasma cytokines, especially TNF- $\alpha$ , IL-1 $\beta$ , and IL-6. TNF- $\alpha$  has been shown to cause changes in neurotransmission of the central nervous system leading to behavioral changes such as lethargy and anorexia (Ryan et al., 2007).

Pain symptom was identified by two characteristics, hand feet and back cramps and abdominal & back lancinating. This complaint is felt by the participants every day and more severe during the activity. Pathophysiology of cancer pain involves somatic, visceral and neuropathy. Somatic pain is usually superficial and arises from the skin, soft tissue and musculoskeletal tissue. Visceral pain is dull originating from internal organs. Neuropathic pain arises from the structure of the somatosensory system such as receptors, peripheral nerves, autonomic nerves and central nervous system (Mishra, Bhatnagar, Chaudhary, & Rana, 2009). Another research reported that 13% of breast cancer patients complained of moderate to severe pain (rating > 4) at early of radiotherapy and 23% reported moderate to severe pain after radiotherapy (Parks et al., 2017). Persistent pain also felt in breast cancer patients after 5-7 years post-chemotherapy (Mejdahl, Andersen, Gärtner, Kroman, & Kehlet, 2013).

Details characteristic of nausea/vomiting were smelling food and at any time and it sometimes followed vomiting, it cause reduce nutritional intake and body weight. Participants only able to spend half the portion of the meal. Vomiting is a reflex triggered by a toxic substance, such as a chemotherapy agent, which causes damage to the gastric mucosa and the small intestine, thus stimulating the vagal afferent interacting with the central nervous system, and generating an emetic response (Hesketh, 2008). Neurotransmitters that affect the emetic response are serotonin, dopamine, acetylcholine, and  $\gamma$ -aminobutyric acid (GABA), histamine, endorphins, and cannabinoids (Navari, 2009). Cancer patients in another research stated that the effects of chemotherapy led to nausea of 77%, vomiting 75% (Aslam et al., 2014).

Insomnia characterized by trouble getting started sleeping & woke up early and woke up at night and could not sleep anymore. It causes participants often feel tired throughout the day. Radiation and chemotherapy are both reported to produce sleep disturbances (Liu & Ancoli-Israel, 2008). In cancer patients, circadian rhythm disturbance is caused by endocrine changes (such as cortisol, melatonin, prolactin), metabolic processes (such as temperature and circulation levels of proteins), and the immune system (such as leukocyte and neutrophil levels) (Liu & Ancoli-Israel, 2008). Sleep disturbances and increased daytime drowsiness occur in adult and child of various types of cancer including acute lymphocytic leukemia, central nervous system tumor, Hodgkin's lymphoma, soft tissue sarcoma, and bone tumor (Mulrooney et al., 2004).

The symptoms of hair loss is suffered in stage 4 of cancer with degrees of hair loss vary, but it does not make the patient feel ashamed. Research conducted on Caucasian cancer patient reported that post-chemotherapy cause hair loss and alopecia on a moderate scale (Kluger et al., 2012).

Constipation is suffered in stage 4 of cancer. Majority of participants told that difficulty defecation with the fastest frequency of defecation is 1 week once times with stiff consistency



of the feces. Constipation often occurs 50-87% in patients with advanced cancer (Abernethy, Wheeler, & Zafar, 2009). Constipation is the third most common symptom in patients receiving cytotoxic chemotherapy with an overall prevalence of 16%, with 5% being severe and 11% moderate (Anthony, 2010). Chemotherapy and radiation drugs affect the contractile physis activity that contribute to colonic motor activity and peristaltic reflexes (Kuizenga et al., 2015).

### **Self-treatment to reduce the symptoms**

The majority of participants who suffered fatigue cope with resting (reduced daily activities and used the time to lie down, while some light activity such as reading, making handwork is done by sitting). But there is a small percentage of participants who overcome fatigue by doing yoga exercise, because they have understood the steps of yoga through social media and training from local cancer organization where they come from. In prior studies reported that the activity of yoga for 12 weeks can reduce fatigue significantly (Bower, Garet, & Sternlieb, 2011).

In pain symptoms, the majority of participants overcome by taking analgesic drug from the physician and combined with the deep breathing relaxation and distraction technique. It has learned from health care workers in the hospital. But sometimes the pain is still felt in moderate to mild scale. Meditation, progressive relaxation, rhythmic respiration, therapeutic touch, transcutaneous electrical nerve stimulation (TENS), hypnosis, music therapy, acupressure are among the non-invasive methods of reducing pain in cancer patients (Woolf, 2011). Non-opioid analgesic drugs are the first step to reduce cancer pain, but if the pain is not well controlled then strong opioids such as morphine can be given (PDQ Supportive and Palliative Care Editorial Board, 2002).

In Nausea/ vomiting symptoms, the majority of participants cope with a combination of antiemetic drug and deep breathing relaxation techniques. In the prior studies reported that providing a relaxation intervention with music for 25 minutes is effective in reducing the intensity of nausea and vomiting in cancer patients (Arakawa, 1997).

In insomnia symptom are overcome by the majority of participants with deep relaxation techniques such as meditation and mild massage on the head/ hands/ feet/ back performed by family caregiver. Medical massage therapy interventions on cancer patients showed significant differences in sleep quality scores before and after intervention in the treatment group (Kashani & Kashani, 2014). Massage therapy can increase different levels of relaxation in a person's body and can improve sleep habits (Reza et al., 2010).

In hair loss symptom, the majority of participants doing nothing of treatment, but there is a small percentage of patients who use hair tonic to cool the head and stimulate hair growth. Scalp cooling especially in the frontal, superior and occipital areas of the head can reduce hair loss during chemotherapy (Ghaffari M, M, AF, & M, 2015).

In constipation symptoms overcome by increasing the high fiber foods consuming such as fruits, vegetables, and many waters, it help them alleviate perceived complaints. Dietary fiber intake increases the frequency of defecation and improves fecal consistency, which is especially useful for patients who are constipated (Yang, Wang, Zhou, & Xu, 2012).




Fig. 2 The Patterns of Symptoms and Self Treatment due to Radiotherapy and Chemotherapy Based on Different Cancer Stage

Fig. 2 shown that side effect of cancer therapy (radiotherapy and chemotherapy) cause the complicated symptoms and its details. Many self- treatment done by patients to reduce the symptoms.

## CONCLUSION

Radiotherapy and chemotherapy in cancer patients can lead to the physical symptoms that disrupt the balance of individual physiological basic needs fulfilled. Every stage of cancer has different details characteristic of symptoms. The Patients did the self-treatment to reduce the symptoms and every stage of cancer has different ways to overcome it.

## REFERENCES

- Abernethy, A. P., Wheeler, J. L., & Zafar, S. Y. (2009). Detailing of gastrointestinal symptoms in cancer patients with advanced disease: new methodologies, new insights, and a proposed approach. *Current Opinion in Supportive and Palliative Care*, 3(1), 41–49. <https://doi.org/10.1097/SPC.0b013e32832531ce>
- Anthony, L. B. (2010). Diarrhea, Constipation, and Obstruction in Cancer Management. In *The MASCC Textbook of Cancer Supportive Care and Survivorship* (pp. 249–260). Boston, MA: Springer US. [https://doi.org/10.1007/978-1-4419-1225-1\\_26](https://doi.org/10.1007/978-1-4419-1225-1_26)
- Arakawa, S. (1997). Relaxation to reduce nausea, vomiting, and anxiety induced by



---

chemotherapy in Japanese patients. *Cancer Nursing*, 20(5), 342–349.

- Areti, A., Yerra, V. G., Naidu, V., & Kumar, A. (2014). Oxidative stress and nerve damage: Role in chemotherapy induced peripheral neuropathy. *Redox Biology*, 2, 289–295. <https://doi.org/10.1016/j.redox.2014.01.006>
- Aslam, M. S., Naveed, S., Ahmed, A., Abbas, Z., Gull, I., & Athar, M. A. (2014). Side Effects of Chemotherapy in Cancer Patients and Evaluation of Patients Opinion about Starvation Based Differential Chemotherapy. *Journal of Cancer Therapy Jour-Nal of Cancer Therapy*, 5(5), 817–822. <https://doi.org/10.4236/jct.2014.58089>
- Badan Penelitian dan Pengembangan Kesehatan. (2018). *Hasil Utama Riskesdas 2018*. Jakarta.
- Baskar, R., Dai, J., Wenlong, N., Yeo, R., & Yeoh, K.-W. (2014). Biological response of cancer cells to radiation treatment. *Frontiers in Molecular Biosciences*, 1. <https://doi.org/10.3389/fmolb.2014.00024>
- Bock, P. R., Hanisch, J., Matthes, H., & Zänker, K. S. (2014). Targeting inflammation in cancer-related-fatigue: a rationale for mistletoe therapy as supportive care in colorectal cancer patients. *Inflammation & Allergy Drug Targets*, 13(2), 105–111.
- Bower, J. E., Garet, D., & Sternlieb, B. (2011). Yoga for persistent fatigue in breast cancer survivors: results of a pilot study. *Evidence-Based Complementary and Alternative Medicine : eCAM*, 2011, 623168. <https://doi.org/10.1155/2011/623168>
- Bunz, F. (2008). *Principles of Cancer Genetics*. Dordrecht: Springer Netherlands. <https://doi.org/10.1007/978-1-4020-6784-6>
- Ghaffari M, S. M., M, G., AF, K., & M, Z. (2015). Site-Dependence Scalp Cooling System to Prevent Hair Loss during Chemotherapy. *Journal of Bioengineering & Biomedical Science*, 5(2), 1–6. <https://doi.org/10.4172/2155-9538.1000158>
- Gilliam, L. A. A., & St. Clair, D. K. (2011). Chemotherapy-Induced Weakness and Fatigue in Skeletal Muscle: The Role of Oxidative Stress. *Antioxidants & Redox Signaling*, 15(9), 2543–2563. <https://doi.org/10.1089/ars.2011.3965>
- Hesketh, P. J. (2008). Chemotherapy-Induced Nausea and Vomiting. *New England Journal of Medicine*, 358(23), 2482–2494. <https://doi.org/10.1056/NEJMra0706547>
- Kashani, F., & Kashani, P. (2014). The effect of massage therapy on the quality of sleep in breast cancer patients. *Iranian Journal of Nursing and Midwifery Research*, 19(2), 113–118.
- Kluger, N., Jacot, W., Frouin, E., Rigau, V., Poujol, S., Dereure, O., ... Bessis, D. (2012). Permanent scalp alopecia related to breast cancer chemotherapy by sequential

fluorouracil/epirubicin/cyclophosphamide (FEC) and docetaxel: a prospective study of 20 patients. *Annals of Oncology*, 23(11), 2879–2884. <https://doi.org/10.1093/annonc/mds095>

Kuizenga, M. H., Sia, T. C., Dodds, K. N., Wiklendt, L., Arkwright, J. W., Thomas, A., ... Costa, M. (2015). Neurally mediated propagating discrete clustered contractions superimposed on myogenic ripples in ex vivo segments of human ileum. *AJP: Gastrointestinal and Liver Physiology*, 308(1), G1–G11. <https://doi.org/10.1152/ajpgi.00230.2014>

Kummer, F., Catuogno, S., Perseus, J. M., Bloch, W., & Baumann, F. T. (2013). Relationship between cancer-related fatigue and physical activity in inpatient cancer rehabilitation. *Anticancer Research*, 33(8), 3415–3422.

Liu, L., & Ancoli-Israel, S. (2008). Sleep Disturbances in Cancer. *Psychiatric Annals*, 38(9), 627–634.

Mejdahl, M. K., Andersen, K. G., Gärtner, R., Kroman, N., & Kehlet, H. (2013). Persistent pain and sensory disturbances after treatment for breast cancer: six year nationwide follow-up study. *BMJ (Clinical Research Ed.)*, 346, f1865.

Mishra, S., Bhatnagar, S., Chaudhary, P., & Rana, S. P. S. (2009). Breakthrough cancer pain: Review of prevalence, characteristics and management. *Indian Journal of Palliative Care*, 15(1), 14. <https://doi.org/10.4103/0973-1075.53506>

Mulrooney, D. A., Mertens, A. C., Neglia, J. P., Green, D. M., Zeltzer, L., & Robison, L. L. (2004). Fatigue and sleep in survivors of childhood cancer: A report from the Childhood Cancer Survivor Study (CCSS). *Journal of Clinical Oncology*, 22(14\_suppl), 8129–8129. <https://doi.org/10.1200/jco.2004.22.90140.8129>

Navari, R. M. (2009). Pharmacological Management of Chemotherapy-Induced Nausea and Vomiting. *Drugs*, 69(5), 515–533. <https://doi.org/10.2165/00003495-200969050-00002>

Parks, J., Yang, X., Zhang, C., Chen, Z., Miller, A. H., Godette, K. D., ... Torres, M. (2017). Predictors of Breast Pain in Breast Cancer Patients One Year After Whole Breast Radiation Therapy. *International Journal of Radiation Oncology, Biology, Physics*, 99, E41. <https://doi.org/10.1016/j.ijrobp.2017.06.689>

PDQ Supportive and Palliative Care Editorial Board. (2002). *Cancer Pain (PDQ®): Health Professional Version. PDQ Cancer Information Summaries*.

Powers, S. K., & Jackson, M. J. (2008). Exercise-Induced Oxidative Stress: Cellular Mechanisms and Impact on Muscle Force Production. *Physiological Reviews*, 88(4), 1243–1276. <https://doi.org/10.1152/physrev.00031.2007>

Reza, H., Kian, N., Pouresmail, Z., Masood, K., Sadat Seyed Bagher, M., & Cheraghi, M. A. (2010). The effect of acupressure on quality of sleep in Iranian elderly nursing

home residents. *Complementary Therapies in Clinical Practice*, 16(2), 81–85.  
<https://doi.org/10.1016/j.ctcp.2009.07.003>

Rossi, L., Stevens, D., Pierga, J.-Y., Lerebours, F., Reyat, F., Robain, M., ... Rouzier, R. (2015). Impact of Adjuvant Chemotherapy on Breast Cancer Survival: A Real-World Population. *PloS One*, 10(7), e0132853.  
<https://doi.org/10.1371/journal.pone.0132853>

Ryan, J. L., Carroll, J. K., Ryan, E. P., Mustian, K. M., Fiscella, K., & Morrow, G. R. (2007). Mechanisms of cancer-related fatigue. *The Oncologist*, 12 Suppl 1(Supplement 1), 22–34. <https://doi.org/10.1634/theoncologist.12-S1-22>

Schnur, J. B., Ouellette, S. C., Bovbjerg, D. H., & Montgomery, G. H. (2009). Breast Cancer Patients' Experience of External-Beam Radiotherapy. *Qualitative Health Research*, 19(5), 668–676. <https://doi.org/10.1177/1049732309334097>

Werdani, Y. D. W. (2017). EFFECT OF MINDFULNESS MEDITATION ON STRESS LEVEL AND COPING MECHANISM IN CANCER PATIENTS. *Folia Medica Indonesiana*, 53(1), 33. <https://doi.org/10.20473/fmi.v53i1.5488>

Woolf, C. J. (2011). Central sensitization: Implications for the diagnosis and treatment of pain. *Pain*, 152(Supplement), S2–S15. <https://doi.org/10.1016/j.pain.2010.09.030>

Yang, J., Wang, H.-P., Zhou, L., & Xu, C.-F. (2012). Effect of dietary fiber on constipation: a meta analysis. *World Journal of Gastroenterology*, 18(48), 7378–7383. <https://doi.org/10.3748/wjg.v18.i48.7378>