

Regional Case Study

Fluctuations of PM_{2.5} and NO_x Concentration and Their Relationship with Meteorological Factors in the Rural Area (Case Study: Puncak Bogor)

Ni Putu Intan Permata Teani^{1*}, Ana Turyanti¹, Ronald Christian Wattimena²,
Yudith Vega Paramitadevi³, Farhan Kurniawan¹

¹Department of Geophysics and Meteorology, IPB University

²Meteorology, Climatology and Geophysics Agency of the Republic of Indonesia

³Environmental Engineering and Management Study Program, College of Vocational Studies, IPB University

*Corresponding Author, email: niputuintang99@gmail.com



Abstract

Rural areas are generally synonymous with better air quality than urban areas. However, if there is an agricultural activity, it has the potential to release nitrogen from the soil, be oxidized and form NO and NO₂ in the air. These two gases encourage the formation of secondary PM_{2.5} particulates in the air of rural areas. This study aims to analyze fluctuations in PM_{2.5} concentration in rural areas, the effect of its precursor NO_x, and the influence of meteorological factors. The location used as a case study is the Cibereum area, Puncak, Bogor Regency. The data used are PM_{2.5}, NO, NO₂, NO_x concentration data (per 3 hours), rainfall, air temperature, wind speed and direction in 2019 and 2020, which were analyzed using correlation and linear regression and the open-air package provided by RStudio. The results showed that the concentration of PM_{2.5} with NO_x as a precursor had a significant positive correlation in 2019 ($r = 0.68$) and 2020 ($r = 0.63$). Cumulative precipitation affects the concentration of PM_{2.5} and NO_x in the air. Meteorological factors have a small correlation value to fluctuations in PM_{2.5} concentration and NO_x concentration except for air temperature ($r = 0.3$).

Keywords: Correlation, meteorological factors, PM_{2.5}, Puncak Bogor, precursor NO_x

1. Introduction

Air pollution is one of the environmental problems that have a severe impact on health. One of the essential air pollutants is PM_{2.5}, a fine particle with a diameter of $< 2.5 \mu\text{m}$. PM_{2.5} can pass through the filtration of nasal hairs until it reaches the end of the respiratory tract with airflow and can accumulate by diffusion, damaging other body parts through air exchange in the lungs (Xing et al., 2016). PM_{2.5} may contain hazardous substances, for example, heavy metals. PM_{2.5} organic components can cause DNA damage and suppress DNA repair; it can also encourage the replication of damaged DNA fragments, triggering carcinogenesis or the process of cancer formation (Xing et al., 2016).

The concentration of PM_{2.5} in the atmosphere was influenced by anthropogenic sources, natural sources, meteorological factors, and chemical processes. In rural areas, anthropogenic emissions are not as much as in urban areas, so particulate fluctuations are more influenced by natural sources,

meteorological factors, and chemical reactions to form secondary particulates. According to Almaraz et al. (2018), NO_x is the primary pollutant in agricultural areas. Nitrogen oxide (NO_x) is one of the secondary particulate precursors for PM_{2.5}. According to Leibensperger et al. (2011), anthropogenic NO_x emission can affect PM concentration in a complex manner depending on the chemical environment of the receptor area. Research Wang et al. (2020) in China found a significant positive correlation between NO_x concentration and PM_{2.5} concentration due to NO_x conversion affecting PM_{2.5} formation. In Indonesia, research on particulate precursors is still rare due to the limited data available. In addition to NO_x, ammonia (NH₃), VOCs, and sulfur oxides (SO_x) are PM_{2.5} precursor compounds (Hodan & Barnard, 2004).

The time required for the precursor compounds to react and form particulates varies greatly and depends on the precursor chemistry and meteorological conditions. According to Hodan and Barnard (2004), the contribution of NO_x to the formation of PM_{2.5} differs greatly depending on atmospheric conditions, including humidity and temperature. According to Wang-Li (2015), most inorganic PM_{2.5} in the atmosphere is secondary and is formed through an acid or base neutralization process involving NO_x and NH₃ as precursors. NO_x reacts with OH to form HNO₃, which then reacts with NH₃ available in the rural atmosphere to form ammonium nitrate (NH₄NO₃). This chemical reaction is highly dependent on temperature, humidity, and the concentration of the precursor. When the temperature is high and the humidity is low, the equilibrium will shift to the gas phase NH₃ and HNO₃. Vice versa, when the temperature is low and the humidity is high, the equilibrium will move to the NH₄NO₃ particle phase. The chemical equation for the formation of secondary particulates involving NO_x is as follows:



The increase in PM_{2.5} concentration in the afternoon and evening can be influenced by local factors, altitude, and weather patterns in the region, including temperature and humidity. The decrease in temperature, especially when there is an inversion of temperature at night, will inhibit particle diffusion and the accumulation of particles on the surface, thereby increasing the PM_{2.5} concentration. Condensation due to high humidity can also increase PM_{2.5} concentrations (Hernandez et al., 2017). In general, air pollution is mainly analyzed in urban areas. In contrast, rural areas are considered clean areas, indicated by pollutant concentrations that do not exceed the applicable ambient air quality standards. However, the air quality in rural areas can also be polluted by certain pollutants related to the potential for emissions from agricultural activities. According to Majra (2011), López-Aparicio (2013), and Wang-Li (2015), the primary pollutant produced by agrarian movements is ammonia (NH₃), which will contribute to the formation of secondary particulates as a result of the chemical reaction shown in equation 1): According to the research of Chang et al. (2020), the concentration of NH₃ in rural areas is around 4.6, 8.3 g/m³, and about 7-8 g/m³ according to Wang et al. (2015b) in rural China.

Puncak area (Bogor Regency) is one of the rural areas in Indonesia which considered to have clean air. However, the Puncak Bogor area is not only agricultural but also a tourist destination that is very busy with tourists, especially during holidays. This will affect the number of motorized vehicles heading to the Puncak area so that they can contribute to sources of NO_x and PM_{2.5} emissions in the region. Based on data from Jasa Marga (2019), the average number of motorized vehicles passing through the Gadog toll gate towards Puncak was 31467 vehicles per working day and 33206 vehicles per holiday during the pre-pandemic period. This study aims to analyze the fluctuation of PM_{2.5} concentrations and precursor pollutant concentrations (NO_x) in the Puncak Bogor area in 2019 and 2020 and to analyze the influence of meteorological factors that contribute to their fluctuations.

2. Methods

This study examines the Puncak area (Bogor Regency) as one of the rural areas in Indonesia. The data used are PM_{2.5}, NO, NO₂, and NO_x, as well as meteorological data in air temperature, humidity, wind direction, speed, and precipitation, with a study period of 2019 and 2020. The 2019 and 2020 periods are used, taking into account the influence of large-scale social restrictions (PSBB) during the COVID-19 pandemic, which is expected to affect fluctuations in vehicles entering the Puncak area, as well as affect air pollutant emissions. The data above is sourced from the BMKG Cibereum Air Pollution Observatory, a collaboration between the Meteorology, Climatology and Geophysics Agency (BMKG) and the National Institute for Environmental Studies (NIES) Japan. The system continuously monitors many air quality parameters consisting of GHGs and air pollutants (CO₂, CH₄, O₃, CO, NO_x, SO₂, aerosol concentrations PM_{2.5}, PM₁₀, Black Carbon, chemical components and flask sampling of air) [2]. Measurement of CO₂ and CH₄ uses G2301 from Picarro, CO using CO-30r from Los Gatos Research, NO_x using Model 42i-TL from Thermo, SO₂ using Model 43i-TLE from Thermo, O₃ using OA-787 from Kimoto Electric, aerosol concentrations (PM_{2.5}, PM₁₀, BC) and chemical components using ACSA-14 from Kimoto Electric, and automatic flask sampler of air from Koshin-RS (Immanuel et al. 2019). The map of the study location is shown in Figure 1. The monitoring location is located in geographical coordinates 06°34'22.5" South Latitude, 106°44'48.8" East Longitude, and at an altitude of 1130 masl. The area around Cibereum is related to tourism activities, in the form of villas for staying, agricultural areas, as well as tourist attractions, especially in the North and East directions, conditions are more open, and there are many buildings. The inn (villa) location is around 35-800 m. Meanwhile, in the South and Southeast directions, there are mixed green areas between agricultural areas, including tea plantations and Taman Safari Indonesia, as well as mountains that are denser in vegetation, with a higher topography (800 masl). Mount Gede-Pangrango is about 7 km away in this direction, affecting the wind patterns around it by sea breeze and land breeze mechanism.




Figure 1. Map of Cibereum BMKG air quality measurement locations (Source: Google Maps)

The relationship between PM_{2.5} and its precursor, in this case. NO_x was carried out by analyzing the fluctuations in the concentration of the two pollutants and NO_x-related compounds, NO and NO₂. Data fluctuations are displayed through diurnal, daily, and monthly graphs. The effect of precipitation is analyzed through PM_{2.5} and NO_x data plots in real-time in the week with the highest and lowest rainfall. This is done to see the impact of accumulated precipitation conditions on fluctuations in pollutant concentrations. The effect of the wind was analyzed using a polar plot diagram.

Analysis of wind direction and speed was carried out to see the dominant wind direction and wind speed blowing in the Puncak Bogor area and associated with fluctuations in pollutant concentrations. The data is divided into two periods: during the day (07:00-19:00 GMT +7) and night (20:00-06:00 GMT +7). This time separation is to see the influence of mountain and valley winds because of the presence of mountains around the location. Based on Hasanah's research (2018), at night, Cibereum/Citeko gets influenced by local winds. Namely, the wind blows from the Southeast (mountain direction) towards the Northwest (lower plains). Data processing using Microsoft Excel and Rstudio application. Fluctuations in pollutant concentration (PM_{2.5}, NO_x, NO, and NO₂) were analyzed statistically concerning meteorological factors (air temperature, relative humidity, precipitation, wind direction, and speed). Statistical analysis was carried out to see the close relationship between the variables used and to pay attention to the presence or absence of a causal relationship between these variables. Calculations of correlation and regression tests were performed using the Minitab 18 software. The correlation test used in this study was Pearson Correlation.

3. Result and Discussion

3.1. PM_{2.5} and NO_x. Concentration Fluctuations

The annual average PM_{2.5} concentration in 2019 (20.8 µg/m³) was higher than in 2020 (14.8 µg/m³). The average concentration of PM_{2.5} in 2019 exceeded the one-year National Ambient Air Quality Standard of 15 µg/m³, by Government Regulation of the Republic of Indonesia No. 22/2021 concerning the National Ambient Air Quality Standard. In 2020, Large-Scale Social Restrictions (PSBB) were imposed in connection with the COVID-19 pandemic so that the number of motorized vehicles passing through the Puncak Bogor area was reduced and reduced particulate emissions into the air. This condition is following the observations of Wang et al. (2020) in Hangzhou, China which showed a decrease in the average concentration of PM_{2.5} in urban and rural areas of Hangzhou, China, during the implementation of the lockdown.

The diurna pattern of PM_{2.5} concentration in 2019 and 2020 in the Puncak Bogor area has a similar movement pattern, but the magnitude is different (Figure 2). This is indicated by the two lines having a unimodal pattern. Namely, one peak concentration at 19:00 WIB of 29.6 µg/m³ in 2019 and 22.5 µg/m³ in 2020, while the lowest concentration was measured at 07: 00 WIB with a value of 12.6 µg/m³ in 2019 and 8.7 µg/m³ in 2020. This pattern is similar to the results of research conducted by Zhao et al. (2009) in the rural area of Beijing, which shows that PM_{2.5} concentration has a unimodal pattern, with the peak concentration occurring at night and the lowest concentration occurring in the morning. Seasonal and diurnal wind patterns heavily influence the fluctuations in PM_{2.5} concentration in rural areas.




Figure 2. The diurnal average concentration of PM_{2.5} in 2019 and 2020

The average annual concentration of NO₂ in the Puncak Bogor area in 2019 was 11.6 mg/m³, and in 2020 it was 11.11 mg/m³. Both values do not exceed the one-year National Ambient Air Quality Standard value of 50 mg/m³. The pattern of diurnal fluctuations in NO₂, NO, and NO_x concentrations in 2019 and 2020 in the Puncak Bogor area looks similar but has different values, especially since the maximum concentration value in 2019 is higher than in 2020 (Figure 3). This difference is caused by restrictions on community activities when entering the COVID-19 pandemic. Around the location of the Cibereum monitoring station, there are many tourism-related activities, such as hotels or villas and Taman Safari Indonesia, which experienced a decrease in visitors during the pandemic, so the number of transportation needed entering the area is also affected, while also reducing pollutant emissions. According to analysis results in several cities such as China, India, and California, NO₂ and NO_x concentrations decreased during the COVID-19 pandemic in 2020 (Zhang et al., 2021; Liu et al., 2021; Misra et al., 2022; He et al., 2021; Gough & Anderson 2022). The three concentrations started to increase at 05:00 WIB, with the concentration of NO reaching its peak first at 08:00 WIB, while the concentration of NO₂ and NO_x at 09:00 WIB. Then the NO concentration decreased, and around 16:00 WIB experienced a slight increase but not more significant than the first peak. NO_x diurnal fluctuations have a pattern of 2 peak concentrations, morning and evening.

Meanwhile, the concentration of NO₂ and NO_x still has a high value during the day and has a second peak at 17:00 WIB with a higher value. This diurnal pattern is similar to the results of a study conducted by Bassani et al. (2021) and Pancholi et al. (2018), which showed that NO₂ and NO concentrations in rural Italy and Jodhpur India had two concentration peaks occurring in the morning and evening, with the second peak of NO being lower. The concentration of NO_x that increases earlier in the morning can be produced by vehicle emissions from the wind and using fertilizers on agricultural land, especially N-fertilizer (Almaraz et al. 2018). Then, the NO available in the atmosphere reacts with ozone to form NO₂ (equation 5). This causes the time lag at the peak of NO and NO₂ concentrations in the morning Gasmí et al. (2017).



During the day, ozone is still available in the atmosphere because solar radiation assisted its formation, so the NO₂ formation reaction still occurs. Therefore, the concentration of NO₂ and NO_x still has a high value during the day, while the concentration of NO decreases. The higher concentrations of NO₂ and NO_x at the second peak resulted from the gradual accumulation of the reaction products between NO and O₃ to form NO₂, even though the radiation began to decrease (Song et al., 2011). At night, when ozone is unavailable in the atmosphere, the concentrations of NO₂ and NO_x decrease drastically. This happens because at night, the formation of O₃ does not occur, and the source of NO emission is also reduced so that the appearance of secondary NO₂ also decreases (Han et al. 2011). The fluctuation of PM_{2.5} concentration compared to its precursor indicates a time lag when the attention of the two pollutants increases and decreases. Figures 4 and 5 show that the NO_x concentration increased first around 05:00 WIB, while the PM_{2.5} concentration increased at 07:00 WIB. Likewise, when the NO_x concentration decreased first around 17:00 WIB, the PM_{2.5} concentration also decreased at 19:00 WIB.

According to Manning et al. (2018), the diurnal PM_{2.5} cycle is driven by the influence of local emissions, meteorological conditions, and secondary PM_{2.5} production. The concentration of PM_{2.5} increases during the day due to the formation of secondary PM_{2.5} that occurs through the chemical reaction of the precursor gas with solar radiation (Li et al., 2015).




Figure 3. Diurnal average concentrations of NO₂, NO, and NO_x (a) in 2019 (b) in 2020

According to Svoboda et al. (2000), $NO + HO_2 \rightarrow NO_2 + OH$, then NO₂ decomposes in the presence of solar radiation into NO and O, and O reacts with O₂ to produce O₃. Data from Cibereum shows that the diurnal concentration of NO_x will be closely related to NO and Ozone. As result of data analysis at Puncak in 2020, the concentration of PM_{2.5} has decreased, while the concentration of NO_x is not different from 2019.




Figure 4. Fluctuations in average daily, diurnal, and monthly PM_{2.5} and NO_x concentrations in 2019




Figure 5. Fluctuations in average daily, diurnal, and monthly average concentrations of PM_{2.5} and NO_x in 2020

The average daily and monthly NO_x concentrations showed an increase in 2020; this is a different condition from the general conditions during the pandemic, namely a decrease in pollutant concentrations, including NO_x (Misra et al., 2021; He et al., 2021; Bartonova et al., 2022). The conditions around the Cibereum monitoring station are very close to community agricultural activities, so there is a possibility of increased agricultural activities during the lockdown period, which affects N emissions from the fertilizers used, and contributes to the formation of NO_x in the air. However, there is a need for further identification regarding this matter. However, there is a significant difference in the diurnal average hourly PM_{2.5} concentration in 2019 and 2020. Based on this, the decrease in PM_{2.5} concentration in 2020 could be due to a reduction in the primary PM_{2.5} emission source in the Puncak Bogor area. As a result of the implementation of the large-scale social restriction. The daily concentration pattern of PM_{2.5} and NO_x tends to increase on weekends; this is related to the characteristics of the Puncak Bogor area as a tourist destination so that the emission of transportation modes of visitors who come every day contributes to the pollutants found in the Puncak Bogor area. Monthly fluctuations in PM_{2.5} and NO_x concentrations in 2019 and 2020 showed a similar pattern, namely when the NO_x concentration increased, accompanied by an increase in PM_{2.5} concentration, and vice versa when the concentration decreased. Both pollutants tend to grow in the dry season and drop in the rainy season; according to Mukhtar et al. (2013), when it rains, the pollutants floating in the air will be carried away along with the rain.

Based on the results of the correlation analysis, the diurnal average concentration of PM_{2.5} with its precursor concentration had a significant positive correlation in 2019 ($r = 0.68$) and 2020 ($r = 0.63$). These results are similar to those of Wang et al. (2020). They found a significant positive correlation between the concentration of PM_{2.5} and the concentration of NO_x in China due to the second conversion of NO_x affecting the formation of PM_{2.5} in China. The p-value generated in 2019 and 2020 is < 0.05 ; it can be said that the diurnal average concentration of NO_x significantly affects the diurnal average concentration of PM_{2.5} in 2019 (Figure 6).




Figure 6. Scatter plot diagram of the diurnal average concentration of PM_{2.5} with NO_x (a) 2019 (b) 2020

3.2. Analysis of the Effect of Meteorological Factors on PM_{2.5} and NO_x Concentrations

According to Manning et al. (2018), the diurnal PM_{2.5} cycle is driven by the influence of local emissions, meteorological conditions, and secondary PM_{2.5} production. The results of the correlation analysis of pollutant concentrations with meteorological factors resulting in a relatively low correlation coefficient (r), the highest for NO_x with temperature ($r = 0.36$), are shown in Table 1. Common r values for PM_{2.5} per 3 hours and NO_x concentrations per minute in the Cibereum Puncak Bogor area show that meteorological factors are not the dominant factor in influencing fluctuations in the concentration of these pollutants. PM_{2.5} and NO_x were positively correlated with air temperature in 2019 and 2020. According to the research of Wang and Ogawa (2015), An increasing temperature can support the reaction of secondary pollutant formation and increase

PM_{2.5}. Increasing humidity encourages particles to grow more prominent and become too heavy so that deposition occurs to the surface so that the measured particulate concentration decreases.

NO_x concentration has small negative correlation with relative humidity (RH) in 2019 and 2020. This is related to ozone which plays a role in the oxidation reaction process of NO₂ formation. Ozone concentration will increase as temperature increases and humidity decreases. The increased ozone concentration triggers a reaction to the formation of NO₂.

Table 1. Correlation coefficient (r) of PM_{2.5} and NO_x concentrations with meteorological parameters

Pollutant	Air Temperature		Relative Humidity		Precipitation		Wind Velocity	
	2019	2020	2019	2020	2019	2020	2019	2020
	PM _{2.5}	0.15	0.10	-0.16	0.03	-0.03	0.00	-0.08
NO _x	0.36	0.31	-0.17	-0.12	-0.02	-0.04	-0.10	-0.15

The effect of precipitation for real-time data does not show a significant correlation in Cibereum. According to Zhao et al. (2020), rainfall has a washing effect depending on the intensity and the mass of pollutants. High rainfall is more effective at cleaning large particles than small particles. Therefore, an analysis was carried out using the period of the highest and lowest precipitation and a study of monthly rainfall.




Figure 7. Fluctuations in PM_{2.5} and NO_x concentrations for one week (a) high rainfall (b) low rainfall in 2019

PM_{2.5} and NO_x concentration data were analyzed in two time periods, namely 23-29 April 2019 (highest precipitation) and 7-13 September 2019 (lowest precipitation) and 15-21 February 2020 (highest precipitation) and 20-26 August 2020 (lowest precipitation) to see real-time fluctuations in PM_{2.5} concentration and its precursor concentration (Figure 7 and Figure 8). The analysis results show that in periods of high precipitation, the attention of PM_{2.5} and NO_x is lower than in periods of low precipitation, both in 2019 and 2020. Fluctuations in the concentrations of PM_{2.5} and NO_x appear large in the dry season, while changes in concentration are minor in the rainy season. In the rainy season, rainfall effectively reduces pollutant concentrations, which is indicated by the low concentration values in the rainy season in both 2019 and 2020. In 2020, the pollutant concentration decreased, and the decrease in PM_{2.5} was more significant than NO_x. This is because the formation of NO₂ continues to occur with the help of sunlight and ozone. Also, around the monitoring location, there are agricultural areas that are active and have the potential to contribute to N emissions that trigger NO_x formation. PM_{2.5} concentration fluctuations, which are very close to NO_x fluctuations in the dry season of 2020, show that in that year, even though it was dry, the concentration was not as high as in 2019 because it was supported by reduced primary particulate emissions from motor vehicles and community activities

in tourist areas. According to Arunanta (2020), the number of cars crossing the Puncak Bogor area in 2020 has decreased by 70-80% as a result of the implementation of Large-Scale Social Restrictions in the Jakarta area and its surroundings. Therefore, it can be indicated that the decrease in PM_{2.5} concentration in 2020 is more due to reduced vehicle emissions as the primary source of PM_{2.5} in the Puncak Bogor area. Although the monitoring station location is in a higher place and not directly adjacent to the main highway, activities in the surrounding tourist sites also contribute.




Figure 8. Fluctuations in PM_{2.5} and NO_x concentrations (a) high rainfall (b) low rainfall in 2020

Monthly rainfall in the Puncak Bogor area in 2019 and 2020 has a monsoon pattern, namely high rainfall at the beginning and end of the year (Figure 10). Based on monthly data, in the dry season, there is an increase in pollutant concentrations and vice versa. Rain has a role in washing the air (rainwash) and dissolving pollutant particles and other gases in the air. The higher the rainfall, the greater its ability to dissolve gas and pollutant particles in the air (Prabowo & Muslim, 2018). Based on the correlation results, the monthly PM_{2.5} concentration and rainfall have a significant negative correlation in 2019 ($r = -0.80$) and 2020 ($r = -0.63$). The monthly concentration of NO_x also has a negative correlation with rainfall in 2019 ($r = -0.53$) and 2020 ($r = -0.43$). The resulting p-value shows that monthly rainfall significantly negatively affects the average concentration of PM_{2.5} with an R² value of 63.58% in 2019 and 39.60% in 2020, while the NO_x concentration has no significant effect (p-value > 0.05) both in 2019 and 2020. It is estimated that other factors influence NO_x fluctuations around Cibereum and require further research.




Figure 9. Monthly fluctuations of PM_{2.5} concentration, NO_x concentration, and rainfall (a) in 2019 (b) in 2020 in Puncak Bogor

3.3. The influence of Wind Direction and Speed on PM_{2.5} and NO_x Concentrations at Bogor Peak

Wind can significantly affect pollutant concentrations. In general, high wind speeds can reduce pollutant concentrations. Wind speed accelerates the spread of pollutants and can also dilute pollutants (Godish, 1977). Figure 11 shows the distribution of PM_{2.5} and NO_x concentrations along with the wind

direction that blows during the day (Figure 11a) and at night (Figure 11b). This polar plot chart is analyzed based on two time periods: daytime (07:00 to 16:00) and night (19:00 to 04:00) in 2019 and 2020.

At noon in 2019, the maximum concentration of PM_{2.5} was measured along with winds from the Northeast (2-3 m/s) (Figure 11a). This is related to settlements and lodging places such as guesthouses, villas, and other tourist destinations in the Northeast Cibereum area. In addition, there are activities engaged in agriculture that have the potential to produce emissions. In addition, the measured PM_{2.5} and NO_x concentrations were lower when the wind was blowing from the west than the east wind. This can be presumed as a result of the western part of the Cibereum area, which is less densely populated.

In 2020, it was seen that the value of PM_{2.5} concentration during the day decreased (Figure 10b). Due to the existence of Large-Scale Social Restrictions *Pembatasan Sosial Berskala Besar (PSBB)* reducing additional sources of pollutants in the Puncak Bogor area, the activities of the surrounding community only influence those pollutant concentrations. The concentration of NO_x increased along with the wind from the North and Northeast (speed 2-3 m/s).




Figure 10. Polar plot of PM_{2.5} and NO_x concentrations in 2019 and 2020 (a) during the day (b) at night

At night in 2019, the maximum concentration of PM_{2.5} was measured along with winds from the north with low speed (1 m/s), Northeast, and the East (3.5 m/s). This is related to several lodging places and community activities compared to other directions. In addition, Liu et al. (2020) stated that under certain geographical conditions and dominant wind directions, an increase in wind speed can increase the concentration of PM_{2.5}.

The dominant wind direction at night comes from the Southeast, which coincides with the position of Mount Gede-Pangrango, so the wind that blows is a mountain wind. Wind characteristics from the mountains are cleaner, so the polar plot is bluer, indicating low pollutant concentrations.

4. Conclusions

The annual average PM_{2.5} concentration in 2019 (20.8 mg/m³) was higher than in 2020 (14.8 mg/m³), and also the concentration of NO₂ in 2019 was 11.6 g/m³, higher than in 2020 (11.1 g/m³) The diurnal fluctuations of PM_{2.5} concentration in 2019 and 2020 have a unimodal pattern, with peak concentrations occurring at night. Meanwhile, fluxes in NO₂, NO, and NO_x concentrations in 2019 and 2020 have a bimodal pattern, with peak concentrations occurring in the morning and evening.

The fluctuation of diurnal PM_{2.5} concentration had a significant positive correlation with NO_x concentration as precursor ($r_{2019} = 0.68$; $r_{2020} = 0.63$). The meteorological factors such as air temperature, relative humidity, and wind speed had little correlations to PM_{2.5} and NO_x concentrations, whether in 2019 or 2020, except for air temperature to NO_x concentration, which had a correlation coefficient value $r = 0.36$. The influence of precipitation was analyzed in specific periods, mainly high precipitation periods and lower precipitation periods. It showed that the concentrations of PM_{2.5} and NO_x are lower in periods with high precipitation, both in 2019 and 2020. The fluctuation of both pollutants concentrations is higher when precipitation is lower, and vice versa.

Influence of wind analyses showed that the dominant wind that blows during the day in the Puncak Bogor area comes from the West and coincides with lower concentrations of PM_{2.5} and NO_x. Concentrations are higher during the day as winds blow from the North, Northeast to East directions, which is associated with valley breezes, carrying pollutants from residential areas and tourist attractions. While at night, the dominant wind is from the Southeast, along with low concentrations of PM_{2.5} and NO_x. This is related to the Southeast Gede Pangrango Mountains, so the wind from the Southeast is assumed to bring clean air masses from the top of the mountain. However, at night the PM_{2.5} concentration is high along with the wind blowing from the Northeast at low speed and East at the higher rate, which is thought to carry pollutants from around the residential areas (resorts and agriculture).

5. Acknowledgment

We thank Indonesian Agency for Metrological, Climatological, and Geophysics (BMKG) and National Institute for Environmental Japan (NIES Japan) for providing primary and secondary data for this journal.

References

- Almaraz, M., E. Bai, Wang, C., Trousdell, J., Conley, S., Faloona, I., Houlton, B.Z., 2018. Agriculture is a major source of NO_x pollution in California. *Science Advances* 4 (1) eaa03477.
- Arunanta, L.N., 2020. Wisata diperketat karena pandemi, volume kendaraan ke Puncak Bogor turun [WWW Document]. URL <https://news.detik.com/berita/d-5179988/wisata-diperketat-karena-pandemi-volume-kendaraan-ke-puncak-bogor-turun>. (accessed 5.7.21).
- Bassani, C., Vichi, F., Esposito, G., Montagnoli, M., Giusto, M., Ianniello, A., 2021. Nitrogen dioxide reductions from satellite and surface observations during covid-19 mitigation in Rome (Italy). *Environmental Science and Pollution Research* 1 – 24.
- Bartonova, A., Colette, A., Zhang, H., Fons, J., Liu, H-Y., Brzezina, J., Chantreux, A., Couvidat, F., Guerreiro, C., Guevara, M., Kuenen, J.J.P., Solberg, S., Super, I., Szanto, I., Tarrason, L., Thornton, A., Gonzalez-Ortiz, A., 2022. The Covid-19 pandemic and environmental stressors in Europe: synergies and interplays, ETC ATNI report 2021/16, Kjeller.
- Carslaw, D., 2019. An open-air manual is an open-source tool for analyzing air pollution data. London: King's College London.
- Chang, Y., Zhang, Y.L., Kawichai, S., Wang, Q., Damme, M.V., Clarisse, L., Prapamontol, P., Lehmann, M.F., 2020. Convergent estimates of biomass burning-derived atmospheric ammonia in Peninsular Southeast Asia. *Atmospheric Chemistry and Physics Discussions* 1-25.
- Chen, D., Liu, X., Lang, J., Zhou, Y., Wei, L., Wang, X., Guo, X., 2017. Estimating the contribution of regional transport to PM_{2.5} air pollution in a rural area on the North China Plain. *Science of Total Environment* 1, 280 – 291.
- Gasmi, K., Aljalal, A., Al-Basheer, W., Abdulahi, M., 2017. Analysis of NO_x, NO, and NO₂ ambient levels as a function of meteorological parameters in Dhahran, Saudi Arabia. *WIT Transactions on Ecology and The Environment* 211, 77 – 86.
- Godish, T., 1977. *Air Quality*. 4th Edition. Boca Raton: Lewis Publisher, CRC Press LLC.
- Gough, W.A., Anderson, V., 2022. Changing air quality and the ozone weekend effect during the COVID-19 Pandemic in Toronto, Ontario. *Climate* 10, 41.
- Grabowski, B., 2016. “p < 0.05” might not mean what you think: American statistical association clarifies p-values. *Journal of the National Cancer Institute* 108, 1 – 9.
- Han, S., Bian, H., Feng, Y., Liu, A., Li, X., Zeng, F., Zhang, X., 2011. Analysis of the relationship between O₃, NO and NO₂ in Tianjin, China. *Aerosol and Air Quality Research*, 11, 128–139.

- Hasanah, A.U., 2018. Fenomena Long Range Transport pada Dispersi Ozon dan NO_x di Wilayah Bogor. IPB University Scientific Repository.
- He, C., Yang, L., Cai, B., Ruan, Q., Hong, S., Wang, Z., 2021. Impacts of the COVID-19 event on the NO_x emissions of key polluting enterprises in China. *Applied Energy* 281, 116042.
- Hernandez, G., Berry, T.A., Wallis, S.L., Poyner, D., 2017. Temperature and humidity effects particulate matter concentrations in a sub-tropical climate during winter. *Proceedings of International Conference of the Environment, Chemistry, and Biology* 102, 41 – 49.
- Hodan, W.M., Barnard, R.W., 2004. Evaluating the Contribution of PM_{2.5} Precursor Gases and Re-entrained Road Emissions to Mobile Source PM_{2.5} Particulate Matter Emissions. Washington: Federal Highway Administration.
- Immanuel, G.S., Rakhman, A., Budianto, B., Boer. R., Ardiansyah, M., Hamid, N., Gumilang, I., Nishihashi, M., Mukai, H., Terao, Y., Hashimoto, S., Osonoi, Y., Nugroho, R., Suwedi, N., Rifai, A., Ihsan, I.M., Sulaiman, A., Gunawan, D., Suharguniyawan, E., Nugraha, M.S., Wattimena, R.C., 2019. Real time air quality monitoring system in three sites (Bogor, Cibereum and Serpong). *International Conference On Tropical Meteorology And Atmospheric Sciences* 303, 012055.
- Leibensperger, E.M., Mickley, L.J., Jacob, D.J., Barrett, S.R.H., 2011. Intercontinental influence of NO_x and CO emissions on particulate matter air quality. *Atmospheric Environment* 45, 3318 – 3324.
- Li, J., Chen, H., Li, Z., Wang, P., Cribb, M., Fan, X., 2015. Low-level temperature inversions affect aerosol condensation nuclei concentrations under different large-scale synoptic circulations. *Advances Atmospheric Science* 32, 898 – 908.
- Liu, J., Lipsitt, J., Jerrett, M., Zhu, Y., 2021. Decreases in near-road NO and NO₂ concentrations during the COVID-19 pandemic in California. *Environmental Science & Technology Letters* 8, 161–167.
- Liu, Z., Shen, L., Yan, C., Du, J., Li, Y., Zhao, H., 2020. Analysis of the influence of precipitation and wind on PM_{2.5} and PM₁₀ in the atmosphere. *Advances in Meteorology* 13, 1 – 15.
- López-Aparicio, S., Guerreiro, C., Viana, M., Reche, C., Querol, X., 2013. Contribution of Agriculture to Air Quality Problems in Cities and Rural Areas in Europe. Netherland: The European Topic Centre on Air Pollution and Climate Change Mitigation (ETC/ACM).
- Majra, J.P., 2011. Air Quality in Rural Areas, Chemistry, Emission Control, Radioactive Pollution, and Indoor Air Quality. London: IntechOpen.
- Manning, M.I., Martin, R.V., Hasenkopf, C., Flasher, J., Li, C., 2018. Diurnal patterns in global refined particulate matter concentration. *Environmental Science & Technology Letters* 5, 687 – 91.
- Misra, P., Takigawa, M., Khatri, P., Dhaka, S.K., Dimri, A.P., Yamaji, K., Kajino, M., Takeuchi, W., Imasu, R., Nitta, K., Patra, P.K., Hayashida, S., 2021. Nitrogen oxides concentration and emission change detection during COVID-19 restrictions in North India. *Scientific Reports* 11, 9800.
- Mukhtar, R., Panjaitan, E.H., Wahyudi, H., Santoso, M., Kurniawati, S., 2013. Komponen kimia PM_{2,5} dan PM₁₀ di udara ambien di Serpong – Tangerang. *Jurnal Ecolab* 7, 1–7.
- Mulyana, E., 2012. Variasi diurnal angin permukaan di lereng Gunung Gede Pangrango. *Jurnal Sains dan Teknologi Modifikasi Cuaca* 13, 77 – 82.
- Nabilah, F., Prasetyo, Y., Sukmono, A., 2017. Analisis pengaruh fenomena el nino dan la nina terhadap curah hujan tahun 1998-2016 menggunakan indikator ONI (oceanic nino index) (Studi Kasus: Provinsi Jawa Barat). *Jurnal Geodesi Undip* 6, 402 – 412.
- Paiman, 2019. Teknik Analisis Korelasi dan Regresi Ilmu-Imu Pertanian. Yogyakarta: UPY Press.
- Pemerintah Republik Indonesia, 2021. Peraturan Pemerintah Republik Indonesia No. 22 Tahun 2021 tentang Pengendalian Pencemaran Udara. Jakarta: Sekretaris Kabinet Republik Indonesia.
- Pancholi, P., Kumar, A., Bikundia, D.S., Chouraisya, C., 2018. An observation of seasonal and diurnal behavior of O₃eNO_x relationships and local/ regional oxidant (OX $\frac{1}{4}$ O₃ þ NO₂) levels at a semi-arid urban site of western India. *Sustainable Environment Research* 28, 79-89.
- Prabowo, K., Muslim, B., 2018. *Penyehatan Udara*. Jakarta: Badan Pengembangan dan Pemberdayaan Sumber Daya Manusia Kesehatan.

- Song, F., Shin, J.Y., Jusino-Atresino, R., Gao, Y., 2011. Relationships among the springtime ground-level NO_x, O₃, and NO₂ in the vicinity of highways on the US East Coast. *Atmospheric Pollution Research* 2, 374 – 383.
- Wang, J., Ogawa, S., 2015. Effects of meteorological conditions on PM_{2.5} concentrations in Nagasaki, Japan. *International Journal of Environmental Research and Public Health* 12, 9089 – 9101.
- Wang, L., Li, M., Yu, S., Chen, X., Li, Z., Zhang, Y., Jiang, L., Xia, Y., Li, J., Liu, W., Li, P., Rosenfeld, D., Seinfeld, J.H., 2020. The unexpected rise of ozone in urban and rural areas, and sulfur dioxide in rural areas during the coronavirus city lockdown in Hangzhou, China: implications for air quality. *Environmental Chemistry Letters* 18, 1713 – 1723.
- Wang-Li, L., 2015. Insights to the forming secondary inorganic PM_{2.5}: current knowledge and future needs. *International Journal of Agricultural and Biological Engineering* 8, 1–13.
- Wang, S., Nan, J., Shi, C., Fu, Q., Gao, S., Wang, D., Cui, H., Saiz-Lopez, A., Zhou, B., 2015. Atmospheric ammonia and its impacts on regional air quality over The Megacity of Shanghai, China. *Scientific Report* 5, 15842.
- Xing, Y.F., Xu, Y.H., Shi, M.H., Lian, Y.X., 2016. The impact of PM_{2.5} on the human respiratory system. *Journal of Thoracic Disease* 8, 69 – 74.
- Zhang, Z., Liu, Y., Liu, H., Hao, A., Zhang, Z., 2022. The impact of lockdown on nitrogen dioxide (NO₂) over Central Asian countries during the COVID-19 pandemic. *Environmental Science and Pollution Research* 29, 18923–18931.
- Zhao, X., Sun, Y., Zhao, C., Jiang, H., 2020. Impact of precipitation with different intensity on PM_{2.5} over specific regions of China. *Atmosphere* 11, 1 – 15.
- Zhao, X., Zhang, X., Xu, X., Xu, J., Meng, W., Pu, W., 2009. Seasonal and diurnal variations of ambient PM_{2.5} concentration in urban and rural environments in Beijing. *Atmospheric Environment* 43, 2893 – 2900.