

Renewable Oil Extracted from Indonesian Srikaya's (*Annona squamosa* sp.) Seed: Another Potent Source for Biodiesel

Masruri,^{1*} Mukesh Sharma,² Warsito,¹ and Prasetyo Adi³

¹Laboratory of Organic Chemistry, Department of Chemistry, Brawijaya University
Jl. Veteran No. 1 Malang, Indonesia 65145

²Research School of Chemistry, Australian National University, Canberra ACT 0200 Australia

³Faculty of Medicine, Brawijaya University, Jl. Veteran No. 1 Malang Indonesia 65145
Email contact : masruri@ub.ac.id

Received 19 April 2012; Accepted 1 July 2012

ABSTRACT

This study looks at the wastes derived from Indonesian fruit as prospect for biofuels. This report investigates the chemical composition of Srikaya (*Annona squamosa* sp.) seed, which is disposed as waste products from traditional markets. The seeds were extracted with various extraction methods and the oil obtained was analysed by means of gas chromatography (GC/FID), gas chromatography-mass spectrometry (GC/MS), infra-red spectrometry and ultra-violet-visible spectrometry. It was found 2 h extraction using soxhlet apparatus with diethyl ether as solvent gave the optimum time extraction. Moreover, five major components were isolated from i.e.: ethyl hexadecanoate, ethyl hexadec-9-enoate, ethyl octadecanoate, 2-hydroxy-1,3-propanediyl hexadecanoate, octadec-9-enaldehyde, and unknown compound, respectively.

Key word: Annonaceae, *Annona squamosa*, Srikaya, FAME, FAEE, biodiesel

INTRODUCTION

Indonesia has a high diversity of indigenous fruits and plants. This become a unique point and able to promote an innovative economic to improve national and local community income. The un-explore of the potency of local fruit remains an issue for increasing their value. With increasing global and national demand for biofuel from plant, it is focused on exploring other new sources of biofuels besides Indonesian palm, since the main source of Indonesian biofuel is the palm oil. oil [1]. Conversion of palm oil's fatty acid into fatty acid ethyl ester (FAEE) and methyl ester (FAME) yields biofuel that can be used directly in diesel motors [2]. For instance, ethyl oleate (FAEE) and methyl oleate (FAME) can be derived from Indonesian crude palm oil (CPO) by transesterification process employing an acid or basic catalyst (Figure 1).




Figure 1. Transesterification of palm oils (R: C16, C18 of aliphatic carbon chain and R': Ethyl or Methyl group)

Srikaya (*Annona squamosa* L.; Javanese known as Srikoyo) [3], is a family plant of annonaceae (Figure 2), and include in this member is sirsat (*Annona muricata*), apel mete (*Annona glabra*), and buah nona (*Annona reticulata*) [5]. East Java Indonesia is the main producer even though it is widely distribution to all the regions. Traditionally is marketed as family fruit for daily consumption or as a source of the other food products in home industry. However, the seeds were disposed as waste.



Figure 2. Srikaya (*Annona squamosal* sp.) fruit [4]

Research toward this fruit especially seed phytochemistry has become interest for its biological activity. Part of this plant was reported had activity as antioxidant [6], insecticide [7], anticancer [8] and antibacterial [9]. Furthermore, the chemical composition from part of plant had been reported as well with different chemical composition according to the procedure isolation and origine of the plant. Volatile constituent mainly contained garmacrene, caryophylene oxide, and kaur-16-ene. These were obtained by steam distillation [9a]. Tetrafuranic acetogenine was also isolated using petroleum eter extraction [10]. Kaur-16-en-18-oic acid was isolated from lipid fraction, and α -pinene, sabinene, limonene including isorecinoleic acid were detected in essential oils of seed [11]. Cyclopeptide squamin-A was also reported composing seed polar extract [12], anti-malarial alkaloide was also reported as N-nitrosoxylophine, roemerolidine and duguevalline [13].

EXPERIMENT

Material and instrumentations

Annona squamosa fruit was bought from traditional local market in Malang, East Java, Indonesia. The sample was identified in the Laboratory of Taxonomy, University of Brawijaya by drs Djati Batoro, and the specimen was deposited in this laboratory. The seeds for this study were collected from ripened fruit, and grinded using home-blender without a prior drying process. The powdered seed was wrapped with Whatman paper for soxhlet extraction.

The chemicals used were analytical grade; diethyl ether (Merck), ethyl acetate (Merck), chloroform (Merck), n-hexane (Merck), precoated TLC silica gel F254 plate (Merck). Gas chromatography (Shimadzu GC-14B, HPS 50 m column, T 200-260 °C, 5 °C/min, helium flow rate 30 mL/min, FID), GCMS (Shimadzu QP-5000, EI 70 eV, CBP-5 50 m column, T 100-280 °C, 10 °C/min, helium flow rate 0.4 mL/min), FTIR (Shimadzu FTIR-8201PC) using thin film on NaCl plate), and UV-Vis spectrophotometer (Shimadzu UV-160A UV-VIS).

Extraction

A 500 mg of srikaya's seed powder was extracted using soxhlet with diethyl ether (150 mL) as a solvent. A 0.1 mL crude sample was taken every hour for 5h for GC/FID monitoring. The extraction was continued for 5 h. After 5 hours the ether extract was dried using MgSO₄, filtered and evaporated in-vacuo to yield a yellowish-brown oils (1.22 mL, d. 0.96 g/mL, 23.4% yield). Analysis of the oil was performed using FTIR, UV-Vis, and GCMS. The mass spectra resulted was mainly used for further chemical identification by comparing the spectra resulted with known spectra on the library provided from National Institute of Standard and Technology (NIST).

RESULT AND DISCUSSION

The progress of soxhlet extraction of the grounded seed was monitored by GC/FID. Aliquotes of 0.1 mL was collected every hour for 5 hours. Figure 3a is displayed GC chromatogram of sample for 2 hour extraction. Minimum of five compounds were present with two major components indicated by peak 3 and 4. Extraction was continued for 5 h, when the extraction process indicated clear appearance. Concentration of the solvent under reduced pressure afforded the yellow-brown oils (23.4% yield, d. 0.96 g/mL, 25 °C). The graph presented in Figure 3b shows the correlation between compositions of each compound/peak over 5 h. It gave optimum extraction at 2 h, at which compound with retention time of 15.96 min was present as a dominant component, however it decrease afterwards and compound with retention time of 15.19 min was found to be dominant at later stages of the extraction. In general, all components were found to be present in highest percentage in sample obtained for the 2 h extraction




Figure 3. Chromatogram of GC/FID after 2 h soxhlet extraction (a), and their composition after 5 h extraction (b)

Determination by GCMS of the oils provided slightly similar chromatogram, and showed the presence of 7 compounds. It was recorded 7 compounds composed the oils (Figure 4). Analysis of the mass spectrum and comparison with the library spectra (NIST databased spectra) resulted in the identification of 6 of the 7 compounds (Table 1). Compound at t_R 31.10, 33.13, and 9.36 minute were known as fatty acid ethyl ester (FAEE). Importantly, this FAEE composed 81.7% of the total srikaya oils.




Figure 4. The GCMS chromatogram of srikaya's renewable oils

Table 1. Composition of the renewable oil from srikaya's seed

No	t_R (minute)	Percentage	Name	Molecular formula (MW)
1	25.24	1.21%	2,6-di- <i>tert</i> -butyl-3,5-bis(3-methylbut-2-en-1-yl)phenol	$C_{24}H_{38}O$ (342)
2	31.10	14.9%	Ethyl hexadecanoate	$C_{18}H_{36}O_2$ (284)
3	33.12	57.4%	Ethyl hexadec-9-enoate	$C_{20}H_{38}O_2$ (310)
4	33.40	9.36%	Ethyl octadecanoate	$C_{20}H_{40}O_2$ (312)
5	34.84	1.32%	2-Hydroxy-1,3-propanediyl hexadecanoate	$C_{35}H_{68}O_5$ (568)
6	37.66	7.67%	Octadec-9-enaldehyde	$C_{18}H_{34}O$ (266)
7	37.92	8.14%	Unknown	$C_{24}H_{38}O$ (342)

Fragmentation analysis of the FAEE spectra supported the determination. Ethyl hexadecanoate spectra (Figure 5) recorded ion molecule at m/z 284. Further fragmentation through McLafferty rearrangement on α and β position resulted peak at m/z 88 and 101 [14]. In the other spectra fragmentation for ethyl 9-octadecenoate (Figure 6) and ethyl octadecanoate (Figure 7) gave a similar pattern cleavage. Both provided ion molecule peaks at m/z 310 and 312, respectively.




Figure 5. Mass spectra t_R 31.10 min, ethyl hexadecanoate (Biodiesel type FAEE)




Figure 6. Mass spectra t_R 33.12 min, ethyl 9-octadecenoate (biodiesel type FAEE)




Figure 7. Mass spectra t_R 33.40 min, ethyl octadecanoate (Biodiesel type FAEE)

In addition to the identified FAEE compounds, one aldehyde compound was detected as octa-9-decenaldehyde (t_R 37.66 min) in 7.67% (Figure 8). Two compounds were also detected as diisoprenylated di-tert-butylphenol and 1,3-disubstituted stearyl bishexadecanoate. Both of these were detected in 1.21% and 1.32%, respectively. Moreover, a compound (t_R 37.92 min) with molecular formula $C_{24}H_{38}O$ and molecular weight 342 remains unidentified.




Figure 8. Mass spectra t_R 37.66 min, octadec-9-enaldehyde




Figure 9. Infrared spectra (a) and UV-Vis spectra (b) of oil extracted from Srikaya's seed

The infrared spectrometer spectra gave clear evidence of the functional group of FAEE composed the srikaya oils (Figure 9a). A strong band at 1745 cm^{-1} indicated carbonyl group stretching absorption. The presence of ester was also supported with peaks in the fingerprint region for C-O-C bending absorption at 1061 and 1047 cm^{-1} . No significant hydroxyl, amine and amide band was detected. The UV-Vis spectra gave λ_{max} for the oils at 275 nm (Figure 9b).

CONCLUSIONS

To summarise, srikaya seed contained a dominant (81.7%) and potential fatty acid ethyl ester (FAEE), a type of recent developed renewable oil for biodiesel. Three structures FAEE found were ethyl hexadecanoate, ethyl 9-octadecenoate and ethyl octadecanoate.

ACKNOWLEDGMENT

MSR thank to PT. POS Indonesia for scholarship, Laboratory of Organic Chemistry Gadjah Mada University for GCMS analysis. Part of this research was submitted to fulfill part of degree requirement in Chemistry.

REFERENCES

- [1] Suratin Subur, Pabrik Biodiesel Terintegrasi: Terobosan untuk Mempercepat Pengembangan Biodiesel, Lembaga Riset Perkembangan Indonesia, http://www.ipard.com/art_perkebun/may24-06_wrs.asp was accessed on 15 December 2006
- [2] (a) Divya Bajpai and V. K. Tyagi, *Journal of Oleo Science*, **2006**, 55 (10), 487-502; (b) Choo Yuen May, Ma Ah Ngan, Cheah Kien Yoo, Rusnani Abdul Majid, Andrew Yap Kian Chung, Harrison Lau Lik Nang, Cheng Sit Foon, Yung Chee Liang, Puah Chiew Wei, Ng Mei Han and Yusof Basiron, *Palm Oil Development*, 42, 3-7 (Downloaded from <http://download/Palm%20Oil%20Developments/june%202005/Palm%20Diesel-%20Green%20and%20Renewable%20Fuel%20From%20Palm%20Oil.pdf>, date access July 30, 2012).
- [3] Heyne, K., Tumbuhan Berguna Indonesia, 1987, Badan Penelitian dan Pengembangan Kehutanan, Departemen Kehutanan, Jakarta ((Indonesian).
- [4] Picture was taken from <http://t1.gstatic.com/images?q=tbn:ANd9GcQkVMDF8F-H3aOOAFsJAFyPUHFgnARTKEv-PRI24f6OqfWlUsXq> (downloaded in May 8th, 2012).
- [5] (a) Steenis, G.G.J, 1978, Flora untuk sekolah di Indonesia, PT Pradnya Paramita, Jakarta; (b) Kapil Vyas, Hansraj Manda, Rahul K. Sharma and Gaurav Sinhal, *Int. J. Pharm. Th.*, **2012**, 3 (2), 107-118
- [6] Panchawat S., Rathore K.S. and Sisodia S.S., *Int. J. Pharm. Tech. Res.*, **2010**, 2 (1), 232-239.
- [7] (a) Amini, Study on nsectisidal principal of seed of annona species found in Indonesia, Project report of University of Gadjah Mada and USAID, May 31, **1988**; (b) S. Anita, P. Sujatha, and P. Prabudhas, *Recent Res. Sci. Technol.*, **2012**, 4 (20), 19-23.
- [8] (a) B. V. V. Pardhasaradhi, Madhurima Reddy, A. Mubarak Ali, A. Leela Kumari and Ashok Khar, *J. Biosci.*, **2005**, 30 (2), 237-244; (b) K. Suresh, S. Manoharan, K. Panjamurthy and K. Kavita, *Pak. J. Biol. Sci.*, **2006**, 9 (14), 2600-2605.
- [9] (a) Chavan M. J., Shinde D. B., and Nirmal S. A., *Nat. Prod. Res.*, **2006**, 20 (8), 754-757; (b) Padhi, L. P., Panda, S. K., Satapathy, S. N. and Dutta, S. K., *J. Agric. Tech.*,

- 2011**, 7 (1), 133-142; (c) P.S. Asher, Y. S. Shinde and P. P. Chavan, *IJPSR*, **2012**, 3 (5), 1457-1460.
- [10] Hiroshi Araya, *Bull. Natl. Inst. Agro-Environ. Sci.*, **2004**, 23, 77-149.
- [11] Eloisa Helena A. Andrade, Maria das Gracas B. Zoghbi, Jose Guilherme S. Maia, Heinz Fabricus, and Friedhelm Marx, *J. Food Compos. Anal.*, **2001**, 14 (2), 227-232.
- [12] Jian Xia SHI, Hou Ming WU, Fa Hu HE, Kenichiro INOUE, and Zhi Da MIN, *Chinese Chem. Lett.*, **1999**, 10 (4), 299-302
- [13] Tylor Johns, Arnold Windust, Thomas Jurgens, and Saleh Muhammad Mansor, *Phytopharmacology*, **2011**, 1 (3), 49-53.
- [14] R. M. Silverstein, X. W. Francis and J. K. David, *Spectrometric Identification of Organic Compounds*, 7th edition, **2005**, John Wiley & Sons Inc., USA