

## Physicochemical and Stability of Goat Cheese with Mono and Mixed Culture of *Lactobacillus plantarum* and *Lactobacillus rhamnosus*

Triana Setyawardani<sup>1</sup>, Winiati Pudji Rahayu<sup>2</sup>, Nurheni Sri Palupi<sup>2</sup>

<sup>1</sup>Department of Animal Production, Faculty of Animal Science, University of Jenderal Soedirman  
Purwokerto 53123, Indonesia

<sup>2</sup>Department of Food Science and Technology, Faculty of Agricultural Engineering and Technology, Bogor Agricultural  
University, Bogor 16002, Indonesia

Corresponding author email: trianaunsoed@gmail.com.

**Abstract.** The purpose of the study is to assess the effect of the use of mono probiotic culture and mixed cultures of *Lactobacillus plantarum* and *Lactobacillus rhamnosus* on the physicochemical properties and its viability during storage. The soft cheeses were made through three treatments: (1) the use of *Lactobacillus rhamnosus* culture, (2) the use of *Lactobacillus plantarum* culture and (3) the use of mixed cultures (*Lactobacillus rhamnosus* and *Lactobacillus plantarum*). The variables measured were the cheese texture (firmness and stickiness), proximate, pH of the product, and the number of LAB. The results showed that cheese firmness ranged from 10.78 to 47.75 gf, cheeses stickiness was -8.23 to -11.53 gs, cheese pH was 4.70- 5.60; number of cheese LAB was 8.59-9.69 log cfu/g. The content of protein, fat, and ash were 13.65-16.54%, 15.28-20.03%, and 2.7-3.39%, respectively. The conclusion of this study was that the soft cheeses with mixed cultures of *L.rhamnosus* and *L.plantarum* are potentially good to be developed as a probiotic food.

**Key words:** Soft Cheese, *Lactobacillus rhamnosus*, *Lactobacillus plantarum*, Culture

**Abstrak.** Penelitian bertujuan untuk mempelajari pengaruh penggunaan probiotik secara tunggal dan campuran dari *Lactobacillus plantarum* dan *Lactobacillus rhamnosus* terhadap sifat fisikokimia dan viabilitas BAL selama penyimpanan. Keju lunak susu kambing terdiri dari tiga perlakuan yaitu (1) menggunakan kultur *Lactobacillus rhamnosus*, (2) menggunakan kultur *Lactobacillus plantarum* dan (3) campuran kultur (*Lactobacillus rhamnosus* and *Lactobacillus plantarum*). Variabel yang diamati adalah tekstur keju, proksimat, pH produk dan jumlah BAL. Hasil menunjukkan bahwa ketegasan/keutuhan keju mempunyai kisaran 10,78–47,75 gf, kelengketan keju adalah -8,23 sampai -11,53 gs, pH keju adalah 4,70–5,60; jumlah BAL keju adalah 8,59-9,69 log cfu/g. Kadar protein, lemak dan abu keju berturut-turut adalah 13,56-16,54%; 15,28-20,03% dan 2,7-3,39 %. Kesimpulan dari penelitian ini adalah bahwa keju lunak dengan kultur campuran dari *L.rhamnosus* dan *L.plantarum* berpotensi baik untuk dikembangkan sebagai makanan probiotik.

**Kata kunci :** Keju lunak, *Lactobacillus rhamnosus*, *Lactobacillus plantarum*, kultur

### Introduction

Cheese as food is believed to have health benefit (Choi et al., 2012) due to bioactive peptide released during protein hydrolysis process and temperature treatment or microbial enzyme (Hafeez et al., 2014), the peptide has many benefits for health including antioxidant, antimicrobial, anti-hypertension and so on (Nagpal et al., 2011). Probiotics Lactic Acid Bacteria (LAB) in cheese will add functional properties of cheese because bacteria are beneficial to health through its role in improving the health of the digestive tract when consumed regularly and fairly. Probiotic bacteria in cheese must stay alive and survive during processing and storage of the product.

Cheese is milk curd coagulated with rennet, separated from the whey and pressed into a solid. Several types of soft cheese produced traditionally made from goat's milk, among other is Batzos produced as the utilization of whey from cheese making originated from Macedonia (Psoni et al., 2003), Camero fresh cheese produced in Spain (Olarte et al., 2001) and Galotry cheese of Greece (Kondyli et al., 2008); Himalayan cheese (Mushtaq et al., 2015).

During processing and storage of cheese, the number of LAB may experience a decrease in the number of the initial culture, but this decrease in the number of LAB is not as fast as other fermentation

products, because the cheese has higher pH, more solid of its consistency and a higher fat content and these condition safe favorable for microbial probiotic (Stanton et al., 1998a). The use of two bacterial cultures in the fermentation process is capable of converting lactose to lactic acid within 3-4 hours, but the single culture takes 12-16 hours (Gurakan et al., 2010). Probiotic cultures such as *Lactobacillus casei*, *Lactobacillus plantarum* and *Lactobacillus rhamnosus* in cheese making has the goal of improving the quality of products and the development of new functional food products.

The purpose of the study was to assess the effect of using single probiotic culture and a mixture of *Lactobacillus plantarum* and *Lactobacillus rhamnosus* on the physicochemical properties of soft goat cheese, and viability of LAB during storage.

## Materials and Method

### Materials and starter culture preparation

Raw milk to manufacture soft goat cheese was purchased from Pegumas Farm (Gumelar, Indonesia), *Lactobacillus rhamnosus* and *Lactobacillus plantarum* isolated from goat milk Ettawa crossbreed (Setyawardani et al., 2011). *Lactobacillus rhamnosus*, and *Lactobacillus* cultures were refreshed in De Man, Rogosa and Sharpe Broth (MRS, Oxoid, UK) media at 37 °C for 24 hours before inoculated. A total of 5% (v/v) of pure culture in MRSB, was added to the goat milk to make starter culture. Milk containing the LAB cultures was incubated at 37 °C for 6 hours.

### Manufacture of goat milk cheese

Goat milk was provided from local goat farm ettawa cross breed as much as soft cheese with probiotic bacteria are selected in vitro and in vivo, namely *L. rhamnosus* and *L. plantarum*. The process of cheese was, according to (Walstra et al., 1999) and (Gardiner et al., 1998) which refers to the manufacturing of cheddar cheese with a few modifications. Modifications made were the filtering process which was immediately after the gel sheating, no cheddaring process, and no pressing process.

Cheese making was started by pasteurizing milk at a temperature of 65°C for 30 minutes, then lowering the temperature to 37°C. The

pasteurized milk was then added by starter culture as much as 5% (v/v) and incubated again at 37°C for 6 hours. The milk was then added by animal rennet as much as 0.06 mL<sup>-1</sup> and incubated at 37°C for two hours to form a gel. Gel was then cut into pieces and allowed to stand for 10-15 minutes. The heating was done at 40°C for 30 minutes. Filtering was applied to separate the whey, leaving a matrix called curd. The separated solids from the whey was added by salt as much as 2% (b/b), then stirred until evenly distributed. The cheese was packed in a container and stored at refrigerator temperature (5°C).

### Cheese texture and chemistry test

Testing texture according to (Buriti et al., 2007) was performed with Universal Testing Machine TA-XT2i type Stable Micro Systems (England). Cheese samples were shaped cylindrically shaped in a diameter of 2.4 cm and height 3.0 cm. The cheese was cut and then placed at room temperature (25°C) for 20 minutes prior to measurement. The variables measured were firmness and cheeses stickiness done using the Texture Expert of Windows 1.20 computer programs. Total protein content was analyzed by micro Kjeldhal method (AOAC, 2006) fat content with soxlet extraction method (AOAC, 2006), and ash content was analyzed with furnace method (AOAC, 2006). The pH value of the sample was analyzed by method (AOAC, 2006) Semisolid or solid samples were taken as much as 10 grams and 10ml of distilled water was added, homogenized, and then the pH was measured using a pHmeter.

### Lactic acid bacteria count

Count of lactic acid bacteria in cheese sample was measured according to (Burns et al., 2008). As much as 20 g sample was put into 180 ml of 2% sterile sodium citrate (b/v), then the sample was homogenized in a stomacher (Seward stomacher 80 Lab blender, UK) for three minutes. Homogen a test was taken as much as 1 ml and diluted up to 10<sup>8</sup>. Samples from the three highest dilutions

were taken as much as 1 ml aseptically and put into a sterile petri dish. The sample was poured with MRSA medium and incubated at 37 °C for 4 hours and each sample was done triplicate.

### Statistical analysis

Data were analyzed by analysis of variance (ANOVA) and a post hoc test of Duncan was performed with SPSS® 17.0 software.

## Results and Discussion

### Cheese texture

The average value of cheese texture (firmness and stickiness) made by the culture of different probiotic LAB are presented in Table 1. The texture of probiotic soft cheese of this research has mean of firmness ranging 10.78-47.75 gf. The firmness of cheese texture varies with the use of different probiotic LAB. The highest mean was the samples of soft cheese with *L. plantarum* probiotic cultures, followed by mixed cultures (*L. plantarum* and *L. rhamnosus*), and the lowest mean was the soft cheese with probiotic *L. rhamnosus*.

Soft texture of the probiotic cheese *L. rhamnosus* was due to the higher water content than other treatments. The water content which is too high in protein matrix results in product which is less elastic and easily broken during the pressing process of cheese. This is because water plays a role as a plasticizer. Water acts indirectly to lower the concentration of casein in the cheese matrix, so that the elasticity is reduced and the product is more easily broken (Fox et al., 2000; Delgado et al., 2011; Magenis et al., 2014). The average water content of soft cheese LAB probiotic *L. rhamnosus* was 60.07%, whereas for probiotics of cheese sample soft *L. plantarum* was 56.42% then followed by LAB mixture (*L. rhamnosus* and *L. plantarum*) of 58.20% (unpublished data). The results showed that the addition of single or mixed cultures resulted in differences in the level of soft cheese stickiness with a range of -

8.23 to-11.53 gs. Cheese obtained with *L. plantarum* treatment was the stickiest among those of the other treatments. Factors affecting the stickiness of cheese are fat and casein which affect the cheese matrix formation (Bryant et al., 1995) argued that the level of cheese stickiness is influenced by fat content which is the higher the fat content, the stickier will the cheese be. Cheese stickiness is one of the characteristics of the texture profiles of cheese which is produced by casein matrix by means of trapping the fat globules. According to (Banks, 1998), the structure of cheese is influenced by the levels of protein, fat and water and biochemical activities during storage.

### Chemical characteristics of goat cheese

Chemical characteristics of goat's milk soft cheese with the addition of different cultures of probiotic LAB are presented in (Table 2). The average of protein content of soft cheese of this research ranged 13.65–16.54%. The use of *L. rhamnosus* culture produced soft cheese with significantly lower protein levels than other treatments. On the other hand, the *L. plantarum* and a mixture culture (*L. rhamnosus* + *L. plantarum*) produced the same protein level.

Table 2 shows that the addition of probiotic cultures treatments affects the fat content of goat milk soft cheeses. The cheese has a fat content of 15.67-20.03%. Addition of *L. plantarum* culture produced the lowest fat content i.e. 15.67 compared to the other treatments. The fat content of this study was higher than the results of (Buriti et al., 2007; Silveira et al., 2015) which is a 15.37%. The ash content in soft cheese with the use of different cultures was significantly different in the treatment of mixed cultures (*L. rhamnosus* and *L. plantarum*) with other treatments and the average ash content was 2.58-3.59%. The highest average of ash content of cheese samples of *L. plantarum* probiotic LAB (3.52%), followed by soft cheeses of *L. rhamnosus* probiotic LAB (3.39%) and the smallest was the

mixture LAB probiotic soft cheese (2.58%). The low levels of ash in the treatment of mixed

Table1. Average value of soft cheese texture using single and mixed cultures

Culture types	Firmness (gf)	Stickiness (gs)
<i>L. rhamnosus</i>	10.78±0.96 <sup>a</sup>	-8.23±0.67 <sup>a</sup>
<i>L.plantarum</i>	47.75±0.79 <sup>c</sup>	-11.53±0.76 <sup>b</sup>
<i>L.rhamnosus+L.plantarum</i>	34.73±1.30 <sup>b</sup>	-10.18±0.73 <sup>ab</sup>

Data values were expressed as means ±SD (n= 6) Means within the same column with different superscripts are significantly different at p<0.05 by Duncan's multiple range test.

Tabel 2. Chemicals composition of soft cheese with single and mixed cultures (%)

Culture types	Protein	Fat	Ash
<i>L. rhamnosus</i>	13.65±0.31 <sup>a</sup>	17.02±1.54 <sup>a</sup>	3.39±0.36 <sup>b</sup>
<i>L.plantarum</i>	15.43±1.28 <sup>b</sup>	15.28±0.75 <sup>a</sup>	3.35±0.29 <sup>b</sup>
<i>L.rhamnosus+L.plantarum</i>	16.54±0.67 <sup>b</sup>	20.03±1.28 <sup>b</sup>	2.58±0.20 <sup>a</sup>

Data values were expressed as means ±SD (n= 6) Means within the same column with different superscripts are significantly different at p<0.05 by Duncan's multiple range test.

cultures (*L. rhamnosus* and *L.plantarum*) was suspected that the mixed cultures require more minerals during cheese making than single culture, consequently minerals remaining in the curd cheese decreased.

### The pH value of cheese

The LAB probiotics activity during processing and storage of the products can be observed by looking at the pH changes and the amount of LAB. The pH data during storage is presented in Figure 1.

Soft cheese storage for four weeks lowered the pH of the product. The decrease in pH begins in the second week to the third week of storage. The average pH during the four weeks of storage was at 4.7-5.6 with the lowest pH observed in the addition of culture *L.plantarum* treatment and the highest was in the addition of probiotic *L.rhamnosus* and mixed cultures. Addition of *L.plantarum* resulted in the lowest pH (5.0) among all treatments (pH 5.7 for *L.rhamnosus* and pH 5.2 for the treatment of mixtures).

The decrease in pH is due to the continuing lactic acid production process and the presence of other organic acids produced by the starter and probiotic cultures (Buriti et al., 2007). The

added LAB cultures will lower the pH of milk by changing the milk lactose into lactic acid. Gel formation of cheese is also affected by pH and temperature. At pH of 5.25 gels formed is solittle because the binding between the micelles are very weak. At low temperatures, the binding between the gel particles is stronger (Walstra et al., 1999). At the end of products storage, the pH increased approaching the initial pH. The increased pH was due to the increased metabolic activities of yeast and molds that utilize lactic acids as the carbon source and or the process of proteolysis by releasing large amounts of alkaline components.

### Lactic acid bacteria count

The highest LAB culture in *L.rhamnosus* curds was 8.64 log cfu/g; 9.83 log cfu/g for *L.plantarum* and 10.04 log cfu g<sup>-1</sup> for mixed cultures. The results of counting the number of LAB during four weeks of storage are presented in (Table 3).

Cheese made with *L.rhamnosus*, *L.plantarum* and mix (*L.rhamnosus* and *L.plantarum*) cultures did not experience a decrease in the number of LAB during four weeks of storage. This shows the well ability of LAB to live on cheese products and the indication of well adaptability of LAB ncheese matrix environment, during the four

weeks. This was caused by the removal of most lactose during whey separation and only little

amount of lactose remained. The increase in lactic acid content during the first few weeks of




Figure 1. The changes of soft cheese pH store for four weeks

Tabel 3. Change of lactic acid bacteria count in soft cheese stored for four weeks (log cfug<sup>-1</sup>)

Cultural types	Ripening period (week)				
	0 <sup>1</sup>	1	2	3	4
<i>L.rhamnosus</i>	8.58±0.13 <sup>a</sup>	8.76±0.07 <sup>a</sup>	8.68±0.18 <sup>a</sup>	8.49±0.09 <sup>a</sup>	8.59±0.14 <sup>a</sup>
<i>L.plantarum</i>	10.04±0.21 <sup>a</sup>	9.88±0.13 <sup>a</sup>	9.93±0.44 <sup>a</sup>	9.89±0.27 <sup>a</sup>	9.87±0.24 <sup>a</sup>
<i>L.rhamnosus</i> + <i>L.plantarum</i>	9.83±0.21 <sup>a</sup>	9.56±0.05 <sup>a</sup>	9.67±0.17 <sup>a</sup>	9.58±0.15 <sup>a</sup>	9.69±0.12 <sup>a</sup>

cfu : colony forming unit, Values are mean ± SD (n = 6).

<sup>1</sup>0 month means the sample obtained after 48 hours from cheese making. Values within the same column with different superscripts are significantly different at p<0.05 by Duncan's multiple range test.

storage is caused by fermentation of lactose remained in the cheese (Ong et al., 2007). The population of LAB in the cheese reduce during processing and storage, although the rate of decrease is lower when compared to other fermented milk. This is due to the high pH of cheese with denser consistency and higher fat content, that provide protection to the LAB during storage and digestion by the gastrointestinal tracts. The population of LAB in the cheese reduce during processing and storage, although the rate of decrease is lower when compared to other fermented milk. This is due to the high pH of cheese with denser consistency and higher fat content, that provide protection to the LAB during storage and digestion by the gastrointestinal tracts (Stanton et al., 1998b)

These microbes should be able to survive the conditions of the stomach and digestive tract so it is useful to maintain the balance of intestinal microbial populations. In cheese making, the number of LAB probiotic cheese is one of the requirements that must be met to obtain functional benefits. It is recommended to consume 10<sup>8</sup>-10<sup>9</sup> cfu in each serving to obtain health benefits (Araújo et al., 2010). The number of LAB trapped in the gel matrix formed after the addition of rennet also reached 10<sup>8</sup> cfu/g so that it has the potential to be developed into probiotic food product.

The study was designed to assess and to evaluate the effect of using single probiotic culture and a mixture of *L.plantarum* and *L.rhamnosus* on the physicochemical properties of goat cheese, and count of LAB during storage. The data from current study indicated that

mixed cultures (*L. rhamnosus* and *L. plantarum*) in soft goats milk cheese has better texture, higher number of LAB, higher protein content, than the culture of *L. rhamnosus*. Soft cheeses made with mixed cultures (*L. rhamnosus* and *L. plantarum*) were potential to be developed as functional cheese.

## Conclusion

Mixed cultures (*L. rhamnosus* and *L. plantarum*) in soft goats milk cheese has better texture, higher number of LAB, higher protein content, than the culture of *L. rhamnosus*. Soft cheeses made with mixed cultures (*L. rhamnosus* and *L. plantarum*) were potential to be developed as functional cheese.

## Acknowledgement

We would like to thank to Dr. RRA Maheswari and Juni Sumarmono PhD for their contribution in the paper corection and preparation.

## References

- AOAC [Association of Official Analytical Chemists]. 2006. Official Method of Analysis. 15<sup>th</sup> Ed. Association of Official Analytical Chemists Inc., Virginia USA.
- Araújo EA, AF de Carvalho, ES Leandro, MM Furtado and CA de Moraes. 2010. Development of a symbiotic cottage cheese added with *Lactobacillus delbrueckii* UFV H2b20 and inulin. *Journal of Functional Foods* 2:85-89.
- Banks JM. 1998. Cheese. In: R. Early (ed.) *The Technology of Dairy Products*. p 81–122. Blackie Academic and Professional, London, UK.
- Bryant A, Z Ustunol and J Steffe. 1995. Texture of Cheddar cheese as influenced by fat reduction. *J. Food Sci.* 60:1216-1219.
- Buriti FC, HR. Cardarelli, TM Filisetti and SM Saad. 2007. Synbiotic potential of fresh cream cheese supplemented with inulin and *Lactobacillus paracasei* in co-culture with *Streptococcus thermophilus*. *Food Chem.* 104:1605-1610.
- Burns P, F Patrignani, D Serrazanetti, G Vinderola, J Reinheimer, R Lanciotti and M Guerzoni. 2008. Probiotic Crescenza Cheese Containing *Lactobacillus casei* and *Lactobacillus acidophilus* Manufactured with High-Pressure Homogenized Milk. *J. Dairy Sci.* 91:500-512.
- Choi J, L Sabikhi, A Hassan and S Anand. 2012. Bioactive peptides in dairy products. *Int J Dairy* 65:1-12.
- Delgado FJ, J González-Crespo, R Cava and R Ramírez. 2011. Proteolysis, texture and colour of a raw goat milk cheese throughout the maturation. *European Food Research and Technology* 233: 483-488.
- Fox T, M Cogan, TP Guinee and PLH McSweeney. 2000. *Fundamental of Cheese Science*. Aspen Publishers. Maryland. USA.
- Gardiner G, R Ross, J Collins, G Fitzgerald and C Stanton. 1998. Development of a Probiotic Cheddar Cheese Containing Human-Derived *Lactobacillus paracasei* Strains. *Applied and environmental microbiology* 64:2192-2199.
- Gurakan G, Candan, Altay and Neslihan. 2010. Yogurt Microbiology and Biochemistry. In: F. Yildiz (ed.) *Development and Manufacture of Yogurt and Other Functional Dairy Products*. p 97-122. CRC Press. Boca Raton. USA.
- Hafeez Z, C Cakir-Kiefer, E Roux, C Perrin, L Miclo and A. Dary-Mouro. 2014. Strategies of producing bioactive peptides from milk proteins to functionalize fermented milk products. *Food Res Int* 63, Part A:71-80.
- Kondyli E, M Katsiari and L Voutsinas. 2008. Chemical and sensory characteristics of Galotyri-type cheese made using different procedures. *Food Control* 19:301-307.
- Magenis RB, ES Prudêncio, CB Fritzen-Freire, MP Stephan, A Silvio do Egito and H Daguier. 2014. Rheological, physicochemical and authenticity assessment of Minas Frescal cheese. *Food Control* 45:22-28.
- Mushtaq M, A Gani, PH Shetty, FA Masoodi and M Ahmad. 2015. Himalayan cheese (Kalari/kradi): Effect of different storage temperatures on its physicochemical, microbiological and antioxidant properties. *LWT - Food Science and Technology* 63:837-845.
- Nagpal R, P Behare, R Rana, A Kumar, M Kumar, S Arora, F Morotta, S Jain and H Yadav. 2011. Bioactive peptides derived from milk proteins and their health beneficial potentials: an update. *Food & function* 2:18-27.
- Olarte C, E Gonzalez-Fandos and S Sanz. 2001. A proposed methodology to determine the sensory quality of a fresh goat's cheese (Cameros cheese): application to cheeses packaged under modified atmospheres. *Food Qual Prefer* 12:163-170.
- Ong L, A Henriksson and NP Shah. 2007. Proteolytic pattern and organic acid profiles of probiotic Cheddar cheese as influenced by probiotic strains of *Lactobacillus acidophilus*, *Lb. paracasei*, *Lb. casei* or *Bifidobacterium* sp. *Int Dairy J.* 17:67-78.

- Psoni L, N Tzanetakis and E Litopoulou-Tzanetaki. 2003. Microbiological characteristics of Batzos, a traditional Greek cheese from raw goat's milk. *Food. microbiology* 20:575-582.
- Setyawardani T, W Rahayu, R Maheswari and N. Palupi. 2011. Identification and characterization of probiotic lactic acid bacteria isolated from indigenous goat milk. *Animal Production* 13.
- Silveira EO, JH Lopes Neto, LAd Silva, AES Raposo, M Magnani and HR Cardarelli. 2015. The effects of inulin combined with oligofructose and goat cheese whey on the physicochemical properties and sensory acceptance of a probiotic chocolate goat dairy beverage. *LWT - Food Science and Technology* 62:445-451.
- Stanton C, G Gardiner, P Lynch, J Collins, G Fitzgerald, and R. Ross. 1998a. Probiotic cheese. *International Dairy Journal* 8:491-496.
- Stanton C, G Gardiner, P Lynch, J Collins, G Fitzgerald and R Ross. 1998b. Probiotic cheese. *Int Dairy J.* 8:491-496.
- Walstra P, TJ Geurtz, A Noomen, A Jellema and MAJS van Boekel. 1999. *Dairy technology: principles of milk properties and processes.* Marcel Dekker Inc. New York. USA.