

# Effect of mangrove leaves extract (*avicennia marina*) concentration on the growth of streptococcus mutans and candida albicans



CrossMark

Mohammad Dharmautama<sup>1</sup>, Richard Tetelepta<sup>1,2\*</sup>, Muhammad Ikbal<sup>3</sup>, Andi EA. Warti<sup>1</sup>

## Abstract

**Objective:** This study aims to determine the effect of mangrove leaves extract (*avicennia marina*) concentration on the growth of *Streptococcus mutans* and *Candida albicans* which are commonly found in removable denture plates.

**Material and Methods:** Production of mangrove leaves extract (*avicennia marina*) is carried out using 96% ethanol solution. Mangrove leaves extract was prepared at concentrations of 2.5%, 5%, 7.5%, and 10% and controlled with denture cleanser materials available in the market. Furthermore, the bacteria activity of *streptococcus mutans* and *candida albicans* (ATCC 10231) were tested, then the data were analyzed by one-way ANOVA and post hoc test (Tukey HSD).

**Results:** Statistical analysis of mangrove leaves extract with 2.5% concentration ( $6.850 \pm 0.127$ ), 5% ( $7.185 \pm 0.007$ ), 7.5% ( $8.360 \pm 0.283$ ), 10% ( $8.485 \pm 0.219$ ), and positive control ( $9.060 \pm 0.226$ ) showed significant inhibition zone differences ( $p < 0.05$ ). The extract with concentration of 7.5% had optimal inhibition growth of *streptococcus mutans* ( $p < 0.05$ ). Mangrove leaves' extract at 10% concentration had not shown inhibitory effects against *candida albicans*.

**Conclusion:** Mangrove leaves extract (*avicennia marina*) is effective for inhibiting the growth of *streptococcus mutans*, but has not shown inhibitory effects against *candida albicans* until it reached a concentration level of 10%.

**Keywords:** *Candida albicans*, Extract, Mangrove, *Streptococcus mutans*

**Cite this Article:** Dharmautama M, Tetelepta R, Ikbal M, Warti, AEA. 2017. Effect of mangrove leaves extract (*avicennia marina*) concentration on the growth of *streptococcus mutans* and *candida albicans*. *Journal of Dentomaxillofacial Science* 2(3): 155-159. DOI: [10.15562/jdmfs.v2i3.436](https://doi.org/10.15562/jdmfs.v2i3.436)

<sup>1</sup>Department of Prosthodontics, Faculty of Dentistry, Hasanuddin University, Makassar, Indonesia

<sup>2</sup>Department of Prosthodontics, Naval Dental Hospital Ladokgi Yos Sudarso, Makassar, Indonesia

<sup>3</sup>Department of Prosthodontics, Dentamedica Care Center, Makassar, Indonesia

## Introduction

Denture stomatitis is often a problem in removable dentures, which is caused by pathogenic microorganisms, particularly *streptococcus mutans* and *candida albicans*, that easily attach themselves to removable denture plates. Denture stomatitis consists of mild inflammation and mucosal erythema that occur in removable denture base, and usually also on maxillary, completely removable denture. This condition usually occurs without symptoms, but when symptoms do appear there are many of them, including burning sensation, bleeding of mucous, and dryness in the mouth. The etiology is however considered multifactorial, including trauma, *candida albicans*, allergies, systemic adverse conditions, surface texture, and permeability of the denture base and the layer of material, which are regarded as some of the main factors causing denture stomatitis. Dentures can generate a number of changes in the ecology of the oral cavity by accumulating microbial plaque on the surface of the denture base.<sup>1</sup>

Another factor that influences biofilm development is the presence of an acquired pellicle (AP) on the denture material surface. The AP is a conditioning film formed immediately after the substratum is exposed to the oral environment by the

selective adsorption of peptides and proteins from the saliva. The presence of a saliva pellicle can alter the substratum properties, such as the surface free energy, because the film composition changes the surface reactivity, which provides different receptor sites for the adherence of microorganisms.<sup>2</sup> At the beginning of the formation of salivary pellicle, gram-positive bacteria *streptococcus sp.* becomes the first bacteria that gets attached to the denture base and forms colonies. One such bacteria is *streptococcus mutans*.<sup>3</sup> *Streptococcus mutans* produces extracellular polysaccharide (PSE) that is not owned by other bacteria. The substrate is an access for bacteria and other fungi to adhere to denture base. Bacteria and fungi will proliferate into plaques. These plaques cause denture stomatitis.<sup>4</sup>

Denture stomatitis treatment involves steps such as oral and denture base hygiene, avoiding the use of denture overnight, putting dentures into substances such as chlorhexidine or sodium hypochlorite and antifungal therapy. While asking how to clean dentures better, it must always be based on the data of patient's earlier dental health and devise methods using that data for maintaining the dental hygiene.<sup>5</sup>

\*Corresponding to: Richard Tetelepta, Department of Prosthodontics, Faculty of Dentistry, Hasanuddin University, Makassar, Indonesia  
[richardtetelepta@gmail.com](mailto:richardtetelepta@gmail.com)

Denture cleaning can be done either mechanically or chemically. Mechanical cleaning is done by using toothbrush and ultrasonic. Chemical cleaning can be done by immersing dentures in disinfectant solutions, alkaline peroxide, alkali hypochlorite, chlorhexidine, sodium hypochlorite, enzymes, and herbs.

The place plants are fertile with various biologically active compounds such as alkaloids, flavonoids, lignins, phenols, sterols, saponins, tannins and terpenes. These metabolites are the major sources of antibacterial, antioxidant and anticancer agents.<sup>6</sup> Plants serve as a reservoir of effective chemotheraputants and provide valuable sources of natural products in control of several bacterial and fungal diseases.<sup>7,8</sup> Antibiotics are the most important weapons to fight against bacterial infections. Many pathogenic organisms are developing plasmid-mediated resistance to popular drugs. Hence, there is a need for the isolation of novel compounds either from microorganisms or from plants. The potential of mangrove plants as a source of new therapeutic agents is still unexplored. The present study is paying attention to the exploitation of leaves of *avicennia marina* L for extraction in different organic solvents to screen for antioxidant activity by ABTS method and antibacterial activity by agar well diffusion method.<sup>9</sup> *Avicennia marina*, *avicennia alba*, *acanthus ilicifolius*, *ceriops decandra* and *excoecaria agallocha* are found to have better potential given their high antimicrobial properties than others. On the other hand, *acanthus ilicifolius* and *avicennia marina* are two mangroves whose leaf extracts were most active in lowest concentrations, indicating the presence of high amount of secondary metabolites and other active components acting synergistically.<sup>10</sup> The ability of mangrove has been proven; therefore, we were interested in determining whether there was an effect of mangrove leaves extract concentration (*avicennia marina*) to inhibit the growth of *streptococcus mutans* and *candida albicans*.

## Material and methods

This research was conducted in September–October 2016. Samples of *avicennia marina*

cleanser (antibacterial and antifungal) on the plate as a positive control. The research was conducted at the Laboratory of Phytochemistry and Microbiology of the Faculty of Pharmacy Hasanuddin University, Makassar Indonesia. This work is an analytical laboratory experimental research. This study used post-test only with control group design.

Mangrove leaves extract (*avicennia marina*) obtained by maceration of 1 kg mangrove leaves inserted into maceration container, soaked in 2 liters of 96% ethanol, stirred and allowed to stand for 3 × 24 hours and then filtered to produce filtrate, which was collected and evaporated with a rotary evaporator to obtain a thick extract of mangrove leaves. Using Minimal Inhibitory Concentration (MIC), 0.25%, 0.5%, 0.75%, 1%, 2.5% and 5% concentrations of mangrove leaves extract (*avicennia marina*) were prepared. Determination of MIC in *streptococcus mutans* was made using SB (sodium broth), the broth was incubated 37°C for 1 × 24 hours; while for *Candida albicans* potato dextrose broth (PDB) was used and the broth was incubated at ±28°C for 2 × 24 hours; observations were made by looking at the turbidity.

To determine the effect of mangrove leaves extract's (at a concentration of 2.5%, 5%, 10%, 15% and a positive control) on *streptococcus mutans* and *candida albicans*, the extract was poured into a paper disc and inserted into a petri dish containing Muller Hilton Agar (MHA) and incubated at 37°C for 1 × 24 hours for *streptococcus mutans* and PDA, and incubated at ± 28°C for 2 × 24 hours for *candida albicans*. The observation made was made by looking at the inhibition zone in the area around the paper disc. Analysis of data used one-way ANOVA statistical tests and Tukey's HSD and SPSS 20.0 version (SPSS Inc., Chicago, IL, USA).

## Results

### MIC Test

The MIC at 5% concentration showed no growth of *streptococcus mutans* [figure 1](#), but there was growth of *candida albicans*, which was identified from the level of turbidity [figure 2](#).

### Activity Test

One-way ANOVA statistical test of results derived for *streptococcus mutans* [table 1](#) showed the smallest zone of inhibition at 2.5% concentration (6.850 ± 0.127 mm) and the largest zone of inhibition was found at 10% concentrations (8.485 ± 0.219 mm) and the positive control was 9.060 ± 0.226 mm [figure 3](#). Differences in inhibition zone between concentration and controls were significant ( $p = 0.000$ ;  $p < 0.05$ ).

The results show that there was no inhibition zone of mangrove leaves extract in *candida albicans*

**Table 1** Differences in the inhibition zones of various concentrations of mangrove leaves extract (avicennia marina) to streptococcus mutans

Intervention	Concentration	n(%)	Inhibition Zone	
			Mean ± SD	p-value
Mangrove leaves	2.5%	2 (20%)	6.850 ± 0.127	0.000*
	5%	2 (20%)	7.185 ± 0.007	
	7.5%	2 (20%)	8.360 ± 0.283	
	10%	2 (20%)	8.485 ± 0.219	
Positive control		2 (20%)	9.060 ± 0.226	
Total		10 (100%)	7.988 ± 0.890	

\*One-way Anova test; p<0,05: significant

**Table 2** Differences in the inhibition zones of various concentrations of mangrove leaves extract (avicennia marina) to candida albicans

Intervention	Concentration	n(%)	Inhibition Zone	
			Mean ± SD	p-value
Mangrove leaves	2.5%	2 (20%)	0.000 ± 0.000	
	5%	2 (20%)	0.000 ± 0.000	
	7.5%	2 (20%)	0.000 ± 0.000	
	10%	2 (20%)	0.000 ± 0.000	
Positive control		2 (20%)	0.000 ± 0.000	
Total		10 (100%)	0.000 ± 0.000	


\*One-way ANOVA test; p < 0.05: significant

**Table 3** Post hoc test of inhibition zones of mangrove leaves extract (avicennia marina) on streptococcus mutans


Intervention (i)	Comparison (j)	Mean Difference (i-j)	p Value
Mangrove leaves 2.5%	Mangrove leaves 5%	-0.335	0.509
	Mangrove leaves 7.5%	-1.510	0.003*
	Mangrove leaves 10%	-1.635	0.002*
	Positive Control	-2.210	0.001*
Mangrove leaves 5%	Mangrove leaves 7.5%	-1.175	0.010*
	Mangrove leaves 10%	-1.300	0.006*
Mangrove leaves 7.5%	Positive Control	-1.875	0.001*
	Mangrove leaves 10%	-0.125	0.963
	Positive Control	-0.700	0.079
Mangrove leaves 10%	Positive Control	-0.575	0.149

until 10% concentration, which also occurred in positive control table 2.


Results of post hoc Tukey's HSD for inhibition zone of streptococcus mutans between each concentration and positive control table 3 also show that there is no significant difference between 2.5% and 5% concentrations, between 7.5% and 10%, and also in 7.5% concentration and in the



**Figure 1** MIC of mangrove leaves extract (avicennia marina) on the growth of streptococcus mutans



**Figure 2** MIC of mangrove leaves extract (avicennia marina) on the growth of candida albicans



**Figure 3** Inhibition zone of mangrove leaves extract (avicennia marina) on streptococcus mutans

positive control. Therefore, 7.5% is considered as the concentration with optimum inhibition effects.

## Discussion

The analysis of this research showed that the highest inhibitory effectiveness of mangrove leaves extract (avicennia marina) to streptococcus mutans

at 2.5% concentration. The contents of the active compound were alkaloids, flavonoids, terpenoids, steroids, and saponins. These compounds are common ingredients used in modern medicine.<sup>11</sup>

Most often ethanolic extracts are considered as more active than aqueous extracts, probably because biologically active substances are better extracted in this solvent. Ethanol is the most commonly used organic solvent, as the finished products are relatively safer to use. Moreover, nearly all of the identified components from plants active against microorganisms are aromatic or saturated organic compounds and most often obtained through initial ethanol or methanol extraction.<sup>12</sup> The ethanol extract of the plant was found to scavenge superoxide and hydroxyl radicals. The extract was also found to inhibit the generation of nitric oxide radical and lipid peroxides. Recent studies have shown that the plant extract has a remarkable hepato-protective effect. The flavonoids present in the plant were found to have hepato-protective and antioxidant capacities.<sup>13</sup>

Owing to the widespread ability of flavonoids to inhibit spore germination of plant pathogens, they have been proposed against fungal pathogens of humans and were shown to possess capacity to inhibit the growth of the opportunistic pathogens such as *Candida albicans*. Flavonoids have bacteriostatic properties, and at higher concentrations flavonoid were found to be able to kill both gram-negative and gram-positive bacteria.<sup>14</sup>

Alkaloids function as either hydrogen-acceptor or hydrogen-donor in hydrogen bonding critical for the interaction between targets; thus, they potentially affect the curative agents on diseases. Alkaloid has antibacterial ability. The mechanism of alkaloid is that it interferes with peptidoglycan that enables the constitution of the components of the bacterial cell, so the cell-wall layers are not fully formed and cause the death of these cells. In alkaloids, there is also a group of nitrogen base that reacts with amino acid compounds that make up the cell walls of bacteria and bacterial DNA. This reaction results in changes in the structure and composition of amino acids, which leads to changes in the genetic balance in the DNA chain; thus, the cell is damaged and promotes bacterial cell lysis, causing the death of bacterial cells.<sup>15</sup>

Terpenoids are also referred as terpenes and are the largest group of compounds. Many terpenes have biological activities that can be used for the treatment of human diseases. Terpenoids have a very broad range of biological activities. To review all the biologically active terpenoids would be a difficult task. Owing to space constraints, we will focus on terpenoids with activities against cancer,

malaria, inflammation, and a variety of infectious diseases (viral and bacterial). Other chemotherapeutic agents for these diseases have been the subject of many excellent reviews.<sup>16</sup>

Tannin causes denaturation of proteins by forming complexes with proteins through the nonspecific power such as hydrogen bonding and hydrophobic effects such as the formation of covalent bonds, inactivation of microbial adhesion to molecule in attaching to host cells, and stimulation of phagocytic cells that play a role in cellular immune response. Tannins not only heal burns and arrest bleeding but also curtail infection while they continue to heal the wound internally. The ability of tannins to form a protective layer over the exposed tissue keeps the wound from getting further infected. Tannins are also beneficial when applied to the mucosal lining of the mouth.<sup>17</sup>

Saponins are soluble in water and insoluble in ether, and like glycosides on hydrolysis, they can also yield aglycones. Saponins are extremely poisonous, as they cause hemolysis of blood and are known to cause cattle poisoning. Saponins work as an antibacterial agent to destabilize the bacterial cell membranes causing lysis of bacterial cell; therefore, the action mechanism of saponin includes the formation of antibacterial group which disrupts the permeability of bacterial cell membrane, resulting in the damage of cell membranes and causing release of various important components in bacterial cells, that is, proteins, nucleic acids, and nucleotides.<sup>18</sup>

In this study, mangrove leaves extract at 2.5% concentration have already provided significant results; out of the all the concentration levels, it is the 7.5% concentration that yielded optimum inhibition of *Streptococcus mutans*.

Unlike inhibition of *Candida albicans*, the results of this study showed no inhibitory zones of mangrove leaf extracts (*Avicennia marina*) on *Candida albicans* up to 10% concentration; it also occurred in positive controls. The *Candida albicans* biofilm cells that grow on denture acrylic are sensitive to be killed by Hst 5 (the N-terminal 24 amino acid segment of Histatin 3 peptide). Surface-coating acrylic with chlorhexidine or Hst 5 effectively inhibits biofilm growth and has potential therapeutic application.<sup>19</sup> This is due to the highly complex mechanisms of *Candida albicans* infection, including adhesion and invasion. Changes include changes in cell morphology/shape from yeast to filament (hifa), biofilm formation, and immune cell evacuation to the host. The ability of *Candida albicans* to adhere to host cells is an important factor in the early stages of colonization and infection. Phenotypic changes to form filaments allow *Candida albicans* to penetrate the epithelium and play a role in infection and spread to host cells. *Candida albicans* can also

form biofilms that are believed to be involved in attacks on host cells and associated with antifungal resistance. Therefore, although the basic components of candida albicans cell wall are similar in yeast and filament forms, surface proteom and number of individuals differ. PAMP present in immune cells differs substantially. In addition, conditions in which the cells grow can cause substantial changes in the cell wall, even if cell morphology is unchanged. Due to its highly ordered and responsive nature, the cell wall is a moving target that poses significant challenges to the host's immune system.<sup>20</sup> It is therefore recommended to do further research on mangrove leaf extract (avicennia marina) with concentration levels exceeding 10%.

## Conclusion

Mangrove leaves extract (avicennia marina) at 7.5% concentration can yield optimum inhibitory effects on the growth of streptococcus mutans. Mangrove leaves extract (avicennia marina) even at 10% concentration cannot inhibit candida albicans.

## Conflict of Interest

The authors report no conflict of interest.

## References

1. Karthikeyan S, Teny F, Deepthi PV. Denture Stomatitis: a brief review. *J Dent & Medical Sci* 2016;15: 114-116.
2. Indira MGC, Wander J da Silva, et al. Influence of substratum position and acquired pellicle on Candida albicans-biofilm. *Braz Oral Res* 2013;27:369-375.
3. Monroy TB, Victor MM, Fernando FM, et al. Candida albicans, staphylococcus aureus and streptococcus mutans colonization in patients wearing dental prosthesis. *Med Oral and Patol Oral Cir Bucal* 2005;1: 27-39.
4. Sato M, Hironori T, MiokoA, et al. Growth inhibition of oral bacteria related to denture stomatitis by anticandidal-chalcones. *Aust Dent J* 1997;42: 343-346.
5. de-Vasconcelos LC, de-Vasconcelos LCS, de-Azevedo ELG, et al. Denture hygiene: importance in denture stomatitis control. *Rev Gaucha Odontol Porto Alegre*; 2013. p. 255-261.
6. Sala A, Recio MD, Giner RM, et al. *J Pharma Pharmacol* 2002;54: 365-371.
7. Khan MR, Kihara M, Omoloso AD. *Natural Product Sci* 2001;7: 9-12.
8. Perez RM. *Pharmaceutic Biolo* 2003;41: 105-157.
9. Nymathullah SMD, Maheswara URV. Antibacterial and antioxidant activity of avicennia marina leaf. *J Chemic Pharmaceutic Res* 2014;6: 252-256.
10. Madhurima B, Punarbasu C. Antimicrobial potential of leaf extracts of ten mangrove species from indian sundarban. *Int J Pharm Bio Sci* 2014;5: 294-304.
11. Agnel RA, Nishanthini A, Mohan VR. Pharmacological characterisation and in vitro antibacterial efficacy of various extract of avicennia marina (forssk) vierh stem (avicenniaceae) against some infectious bacteria used. *IJUPBS* 2013;2.
12. Guntra K, Linda R. Influence of plant extracts on the growth of oral pathogens Streptococcus mutans and candida albicans in vitro. *Proceedings of the Estonian Academy Sci* 2015;64, 1, 62-67.
13. Ganesh S, Vennila JJ. Screening for antimicrobial activity in acanthus ilicifolius. *Arch ApplSci Res* 2010;2: 311-315.
14. Cushnie TTP, Anrew JL. Antimicrobial activity of flavonoids. *Int J Antimicrobial Agents* 2005;26: 343-356.
15. Nyambuya. Alkaloid extracts from combretumzeyheri inhibit the growth of mycobacterium smegmatis. *BMC Complementary and Alternative Medicine* 2017;17: 124.
16. Guangyi W, Weiping T, Robert RB. Terpenoids as therapeutic drugs and pharmaceutical agents. *Natural Products*; 2005. p. 197-227.
17. Praveen KA, Kumud U. Tannins are as tringent. *J Pharmacogn and Phytochem* 2012;1.
18. Charishma Y, Thiruvengadarajan VS. A review on identification strategy of phyto constituents present in herbal plants. *Int J Res Pharm Sci* 2013;4: 123-140.
19. Pusateri CR, Monaco EA, Edgerton M. *Archives of oral biology* 2009;54: 588-594.
20. Neil ARG, Frank L van de Verdonk, et al. Candida albicans morphogenesis and host defence: discriminating invasion from colonization. *Nat Rev Microbiol* 2013;10: 112-122.



This work is licensed under a Creative Commons Attribution