

<http://heanoti.com/index.php/hn>

RESEARCH ARTICLE

URL of this article: <http://heanoti.com/index.php/hn/article/view/hn20227>

Effect of Probiotic Milk containing *L. casei* Shirota on Immunoglobulin G Levels among Anemic Adolescent Girls

Elya Sugianti^{1(CA)}, Annis Catur Adi², Ikeu Ekayanti³, Trias Mahmudiono⁴, Widati Fatmaningrum⁵, Siti Nur Husnul Yusmiati⁶^{1(CA)}Faculty of Public Health, Airlangga University; Indonesia; sugiantielya@gmail.com (Corresponding Author)²Faculty of Public Health, Airlangga University, Indonesia; annis_catur@yahoo.com³Faculty of Human Ecology, Bogor Agricultural Institute, Indonesia; ikeu.ekayanti@gmail.com⁴Faculty of Public Health, Airlangga University, Indonesia⁵Departement of Public Health-Preventive Medicine, Faculty of Medicine, Airlangga University; Indonesia⁶Faculty of Health Sains, Ma'arif Hasyim Latif University, Indonesia

ABSTRACT

Probiotic play important role on immune systems. This study was to investigate effect of probiotic milk containing *L. casei* Shirota on immunoglobulin G levels among anemic adolescent girls. A total of 22 adolescent girls aged 13-18 years with anemia were included on this study. The subjects were separated into 2 groups i.e. a probiotic group and a control group by simple random sampling. During 4 weeks, a probiotic group received 1 tablet of 60 Fe elemental + 250 µg folate acid once a week and probiotic milk once daily. While a control group received 1 tablet of 60 Fe elemental + 250 µg folate acid once a week. Blood samples were collected at 0 and 4 weeks and assayed for immunoglobulin G levels by ELISA. Immunoglobulin G levels were not significantly increase after intervention. Compared with control group, probiotic milk administration in probiotic group did not significantly increased immunoglobulin G levels ($p>0.05$). The probiotic milk containing *L. casei* Shirota did not improve immunoglobulin G levels on anemic adolescent girls.

Keywords: Probiotic, Immunoglobulin G levels, Anemia, Adolescent girls

INTRODUCTION

Anemia is the major serious health problems among adolescent girls⁽¹⁾. Risk of anemia on adolescent due to period of rapid growth and higher requirement as result of menstruation cycle for a month⁽²⁾. In addition, bad food pattern worsen high risk of anemia among adolescent girls⁽¹⁾. Anemia can cause a decrease in the immune system. Previous studies had found the declining of immune system on individu with anemia⁽³⁻⁷⁾.

Probiotic are bacteria which, when administered in adequate amounts, can to promote health benefit on humans⁽⁸⁾. One of health benefit as result of consuming probiotic was the increasing of immunoglobulin levels, especially immunoglobulin G levels. Previous studies had found the effect of probiotic administration on immunoglobulin G levels, but the results were controversial⁽⁹⁻¹⁸⁾. Clinical trials were needed to clearly investigate the effect of probiotic bacteria on immunoglobulin G levels.

L. casei Shirota was probiotic strain can improve gastrointestinal tract. Previous clinical studies had found the effect of *L. casei* shirota on immune systems by inducing levels of NK cells production⁽¹⁹⁾, inducing IL-2 and IFN γ production⁽²⁰⁾, activation on T cells⁽¹⁷⁾, and inducing immunoglobulin secretion⁽²¹⁻²²⁾. However, effect of *L. casei* Shirota on immunoglobulin G levels has not been established. Therefore, the study was to investigate effect of probiotic milk containing *L. casei* Shirota on immunoglobulin G levels among anemic adolescent girls.

METHODS

This study was undertaken at Jabon's Community Health Centers in Sidoarjo District during Agustus 2017 until Oktober 2017. All of the participant had written informed consent before recruitment this study. This study protocol was approved by the Health Research Ethics Committee, Faculty of Public Health Airlangga University

(No. 418-KEPK). This study was double blind, randomized, pretest posttest control group design. A total of 22 adolescent girls aged 13-18 years who had low hemoglobin levels (Hb < 12 g/dl) were taken as subjects. The subjects were separated into 2 groups by simple random sampling, i.e. a probiotic group and a control group. During 4 weeks, a probiotic group received 1 tablet of 60 Fe elemental+250 µg folate acid once a week and 65 ml portion probiotic milk containing 6.5 x 10⁹ cfu/ml *L.casei* Shirota once daily. While a control group received only 1 tablet of 60 Fe elemental+250 µg folate acid once a week. Data of baseline characteristic were obtained by interview. Data of weight and height body were obtained by anthropometric measurement. Blood samples were collected at 0 and 4 weeks and assayed for immunoglobulin G levels by ELISA. data was analysed by independent t test with confidence interval of 95%.

RESULTS

Baseline characteristics of anemic adolescent girls were based on age distribution, family size, maternal education, maternal occupation, BMI-for-age, and frequency of morbidity 1 month before study (Table 1). Subjects were higher on early adolescent age with small family size in both group. Subjects had mother with high levels of education and worked as a housewife in both group. Base on BMI-for-age, subjects had normal nutrition status. Compared with control group, probiotic group had higher frequency of sick 1 month before study.

There were no significant differences between probiotic and control group at baseline characteristics (p>0.05). Similarly, immunoglobulin G levels were similar between probiotic and control group at baseline (p>0.05).

Subjects had high compliance and similar in both group. On average, total compliance score of supplementation Fe+folat was 93.2% in both group. While total compliance score of probiotic milk drink was 95.8% in probiotic group (data not shown).

Table 1. Baseline characteristics of anemic adolescent girls

Characteristics	Group			
	Probiotic (n=11)		Control (n=11)	
	n	%	n	%
Age groups (years)				
• 13-15	7	63.6	7	63.6
• 16-18	4	36.4	4	36.4
Family size				
• ≤ 4	9	81.8	8	72.7
• >4	2	1.2	3	27.3
Maternal education				
• Low	5	45.5	5	45.5
• High	6	54.5	6	54.5
Maternal occupation				
• Proffesion	2	18.2	2	18.2
• Skilled worker	3	27.3	4	36.4
• Housewife	5	45.5	5	45.5
• Others	1	9.1	0	0.0
BMI-for-age-groups				
• Normal	10	90.9	11	100.0
• Overweight	1	9.1	0	0.0
Frequency of sick 1 month before study				
• ≤ 1				
• 2-3	1	9.1	0	0.0
• >3	3	27.3	6	54.5
	7	63.6	5	45.5

No significant between probiotic and control group were identified by mann whitney test, where p>0.05

Tabel 2 shows the average of immunoglobulin G levels before and after intervention in both group. On average, immunoglobulin G levels after intervention showed an increase in both group, but not significant. Compared with control group, probiotic group had higher increases immunoglobulin G levels.

Table 2. Immunoglobulin G levels before and after the treatment in both group

Variable	Group		p-value
	Probiotic	Control	
Immunoglobulin G ($\mu\text{g/ml}$)			
Baseline	64.8 \pm 9.9	66.2 \pm 6.6	0.688 ^b
Endline	85.8 \pm 40.9	76.6 \pm 14.3	0.487 ^b
Change	21.1 \pm 38.4	10.3 \pm 15.5	
p-value	0.099 ^a	0.051 ^a	

^apaired t test, with $p < 0.05$

^bindependent t test, with $p < 0.05$

DISCUSSION

This present study investigated the effect of probiotic milk containing *L. casei* Shirota on immunoglobulin G levels among anemic adolescent girls. There were no increases significantly immunoglobulin G levels as a result consuming probiotic milk containing *L. casei* Shirota.

Similar to other clinical trials^{(9-12),(14),(17)}, present study found that probiotic milk had no effect on immunoglobulin G levels. However, contrast to previous studies, probiotic administration had significantly increased immunoglobulin G levels^{(13),(15-16),(18)}. The present study strongly contrast with animal studies that clearly explained the effects of probiotic on immunoglobulin G levels⁽²³⁻²⁵⁾.

The lack of the effect of probiotic milk containing *L. casei* Shirota on immunoglobulin G levels can be explained by several reasons. The effect of probiotic milk on immunoglobulin G levels may be specific bacteria strain dependent. Other studies with different strains had a higher effect^{(13),(15-16),(18)}. Furthermore, the effect of probiotic may be dose dependent⁽²⁶⁾. Other clinical trials with higher doses and combined several probiotic strains had the effect on immunoglobulin G levels^{(15),(25)}. In addition, the selection of intervention period may be important. previous studies with longer intervention period had significantly increased immunoglobulin G levels⁽²¹⁾.

Present study shows that no effect of probiotic milk on immunoglobulin G levels due to the selection of probiotic strains used, dose and intervention period. However, maybe there were others factors that becomes causes no effect of probiotic milk administration on immunoglobulin g levels. One such factor was the possibility of intestinal microflora under normal conditions so that the immune system was stable conditions⁽⁹⁾. This condition had caused probiotic milk administration becomes undetectable.

CONCLUSION

The probiotic milk containing *L. casei* Shirota did not improve immunoglobulin G levels on anemic adolescent girls. Future study are needed by increasing of dose levels, selecting of specific strain used and selecting of treatment period, and recruiting of participants with low immune systems.

REFERENCES

1. Irianto K. Balanced Nutrition on Reproductive Health (Gizi Seimbang dalam Kesehatan Reproduksi). Bandung: Alfabeta; 2014.
2. World Health Organization. Nutrition in Adolescence-Issues and Challenges for the Health Sector, Issues in Adolescent Health and Development. Newyork; 2005.
3. Ekiz C, Agaoglu L, Karakas Z, Gurel N, Yalcin I. The Effect of Iron Deficiency Anemia on the Function of the Immune System. The Hematology Journal. 2005;5:579-583.
4. Tang YM, Chen XZ, Li GR, Zhou RH, Ning H, Yan H. Effects of Iron Deficiency Anemia on Immunity and Infectious Disease in Pregnant Women. Wei Sheng Yan Jiu. 2006;35(1):79-81.
5. Mullick S, Rusia U, Sikka M, Faridi MA. Impact of Iron Deficiency Anemia on T Lymphocytes and Their Subsets in Children. Indian Journal of Medical Research. 2006;124:647-654.
6. Attia MA, Essa SA, Nosai NA, Amin AM, El-agamy OA. Effect of Iron Deficiency Anemia and Its Treatment on Cell Mediated Immunity. Indian Journal Hematology and Blood Transfusion. 2009;25(2):70-77.
7. Hassan TH, Badr MA, Karam NA, Zkaria M, El Saadany HF, Rahman DMA, Shahbah DA, Al Morshedy SM, Fathy M, Esh AMH, Selim AM. Impact of Iron Deficiency Anemia on the Function of the Immune System in Children. Medicine. 2016;95(47).
8. Mizock BA. Probiotics. Disease-a-Month. 2015;61:259-290.

9. Spanhaak SR, Havenaar G, Schaafsma. The Effect of Consumption of Milk Fermented by *Lactobacillus casei* Strain Shirota on the Intestinal Microflora and Immune Parameters in Humans. *European Journal Clinical Nutrition*. 1998;52:899-907.
10. Marcos A, Warnberg J, Nova E, Gomez S, Alvarez A, Alvarez R, Mateos JA, Cobo JM. The Effect of Milk Fermented by Yogurt Cultures Plus *Lactobacillus casei* DN-114001 on the Immune Response of Subjects under Academic Examination Stress. *European Journal Nutrition*. 2004;43:381-389.
11. Christensen HR, Larsen CN, Kaestel P, Rosholm LB, Sternberg C, Michaelsen KF, Frokiaer H. Immunomodulating Potential of Supplementation with Probiotics: A Dose-response Study in Healthy Young Adults. *FEMS Immunology Medicines Microbiology*. 2004;47:380-390.
12. Olivares M, Diaz-Ropero MP, Gomez N, Lara-Villoslada F, Sierra S, Maldonado JA, Martin R, Rodriguez JM, Xaus J, The Consumption of Two New Probiotic Strains, *Lactobacillus gasseri* CECT 5714 and *Lactobacillus coryniformis* CECT 5711, Boosts the Immune System of Healthy Humans. *International Microbiology*. 2006;9:47-52.
13. Kim HS, Park H, Cho IY, Paik HD, Park E. Dietary Supplementation of Probiotic *Bacillus polyfermenticus*, Bispan Strain, Modulates Natural Killer Cell and t Cel Subset Populations and Immunoglobulin G Levels in Human Subjects. *Journal of Medicinal Food*. 2006;9(3),321-327
14. Giovannini M, Agostoni C, Riva E, Salvini F, Ruscitto A, Zuccotti GV, Radaelli G. A Randomized Prospective Double Blind Controlled Trial on Effects of Long-term Consumption of Fermented Milk Containing *Lactobacillus casei* in Pre-school Children with Allergic sthma and or Rhinitis. *Pediatric Research*. 2007;62,215-220.
15. Alberda C, Gramlich L, Meddings J, Field C, Mc Cargar L, Kutsogiannis D, Fedorak R, Madsen K. Effects of Probiotic Therapy in Critically Ill Patients: A Randomized, Double Blind, Placebo-controlled Trial. *American Journal Clinical Nutrition*. 2007;85:816-823.
16. Paineau D, Carcano D, Leyer G, Darquy S, Alyanakian, MA, Simoneau G, Bergmann JF, Brassart D, Bornet F, Ouwehand AC. Effects of Seven Potential Probiotic Strains on Specific Immune Responses in Healthy Adults: A Double-blind, Randomized, Controlled Trial. *FEMS Immunology and Medical Microbiology*. 2008;53:107-113.
17. Harbige LS, Pinto E, Allgrove J, Thomas LV. Immune Response of Healthy Adults to The Ingested Probiotic *Lactobacillus casei* Shirota. *Scandinavian Journal of Immunology*. 2016;84:353-364.
18. Lee A, Lee YJ, Yoo HJ, Kim M, Chang Y, Lee DS, Lee JH, Consumption of Dairy Yogurt Containing *Lactobacillus paracasei* ssp. *Paracasei*, *Bifidobacterium animalis* sp. *Lactis* and Heat-treated *Lactobacillus plantarum* Improves Immune Function Including Natural Killer Cell Activity. *Nutrients*. 2017;9.
19. Dong H, Rowland I, Thomas LV, Yaqoob P. Immunomodulatory Effects of a Probiotic Drink Containing *Lactobacillus casei* Shirota in Healthy Older Volunteers. *European Journal of Nutrition*. 2013;52(8):1853-1863.
20. You J, Dong H, Mann ER, Knight SC, Yaqoob P. Probiotic Modulation of Dendritic Cell Function is Influenced by Ageing. *Immunobiology*. 2014;219:138-148.
21. Ivory K, Chambers SJ, Pin C, Prieto E, Arques JL, Nicoletti C. Oral Delivery of *Lactobacillus casei* Shirota Modifies Allergen-induced Immune Responses in Allergic Rhinitis. *Clinical and Experimental Allergy*. 2008;38:1282-1289.
22. Gleeson M, Bishop NC, Oliveira M, Tauler P. Daily Probiotic's (*Lactobacillus casei* Shirota) Reduction of Infection Incidence in Athletes. *International Journal of Sport Nutrition Exercise Metabolism*. 2011;21(1):55-64.
23. de Waard R, Garssen J, Snel J, Bokken GCAM, Sako T, Huis JHJ, Veld T, Vos JG. Enhanced Antigen-specific Delayed-type Hypersensitivity and Immunoglobulin G2b Responses After Oral Administration of Viable *Lactobacillus casei* YIT9029 in Wistar and Brown Norway Rats. *Clinical and Diagnostic Laboratory Immunology*. 2001;8:762-767.
24. Sebastian AP, Keerthi TR, Immunomodulatory Effect of Probiotic Strain *Bacillus subtilis* MBTU PBBMI Spores in Balb/c Mice. *International Journal of Engineering and Technical Research*. 2014;2(11):258-260.
25. Karamese M, Aydin H, Sengul E, Gelen V, Sevim C, Ustek D, Karakus E. The Immunostimulatory Effect of Lactic Acid Bacteria in a Rat Model. *Iran Journal Immunology*. 2016;13:220-228.
26. Hatakka K, Savilahti E, Ponka A, Meurman JH, Poussa T, Nase L, Saxelin M, Korpela R. Effect of Long Term Consumption of Probiotic Milk on Infections in Children Attending Day Care Centres: Double Blind, Randomised Trial. *British Medical Journal*. 2001;322,1327-1329.